





LIBRARY OF CONGRESS.

Chap. .... Copyright No. ....

Shelf RJ511  
F73


UNITED STATES OF AMERICA.











# SKIN DISEASES

OF

## CHILDREN

BY

GEORGE HENRY FOX, A.M., M.D.

Clinical Professor of Diseases of the Skin, College of Physicians and Surgeons,  
New York; Physician to the New York Skin and Cancer Hospital, etc.

---

WITH TWELVE PHOTOGRAVURE AND CHROMOGRAPHIC PLATES,  
AND SIXTY ILLUSTRATIONS IN THE TEXT



TWO COPIES RECEIVED

61594032

NEW YORK

WILLIAM WOOD & COMPANY

1897

RJ 511  
F75  
~~RX 11~~  
~~F75~~

COPYRIGHT, BY  
WILLIAM WOOD & COMPANY,  
1897.

## PREFACE.

---

A SERIES of illustrated articles on certain skin diseases which are apt to occur in infancy and childhood appeared in the *American Journal of Obstetrics and Diseases of Women and Children* a year ago, and are now published in book form. While some additions have been made to most of them, the work is far from being a complete description of all the cutaneous affections which occur in childhood. It is merely intended to serve as a concise and practical sketch of the symptomatology and treatment of those eruptions which are most likely to affect the juvenile patients of every family physician.

To Dr. Horace S. Stokes the thanks of the author are due for valuable assistance, especially in the compilation of the appended formulary.

18 EAST 31ST STREET, NEW YORK,  
June, 1897.



## CONTENTS.

---

	PAGE
Alopecia Areata, . . . . .	1
Ringworm and Favus, . . . . .	11
Contagious Impetigo, . . . . .	21
Psoriasis, . . . . .	27
Ichthyosis, . . . . .	35
Eczema, . . . . .	41
Papilloma Lineare, . . . . .	51
Nævi, Pigmented and Hairy, . . . . .	55
Vascular Nævus, . . . . .	61
Lupus and other Tuberculides, . . . . .	67
Lichen Ruber and Lichen Planus, . . . . .	75
Syphilis Hæreditaria, . . . . .	83
Erythema Nodosum, . . . . .	84
Purpura, . . . . .	85
Molluscum, . . . . .	87
Keratosis Follicularis, . . . . .	89
Keloid, . . . . .	90
Scabies, . . . . .	92
Formulary, . . . . .	95
Index, . . . . .	161





# LIST OF ILLUSTRATIONS.

## PLATES.

	FACING PAGE.
Alopecia areata, . . . . .	1
Trichophytosis capitis, . . . . .	11
Impetigo contagiosa, . . . . .	21
Psoriasis, . . . . .	27
Ichthyosis, . . . . .	35
Eczema, . . . . .	41
Papilloma lineare, . . . . .	51
Nævus pilosus, . . . . .	55
Vascular nævus, . . . . .	61
Lupus vulgaris, . . . . .	67
Lichen ruber, . . . . .	75
Syphilis hæreditaria, . . . . .	83

## FIGURES.

FIG.	PAGE.
1. Bald ringworm, . . . . .	2
2. Serpiginous form of alopecia areata, . . . . .	3
3. Alopecia areata—growth of white hair on patches, . . . . .	4
4. Alopecia areata—complete loss of hair, . . . . .	5
5. Trichophytosis corporis, . . . . .	12
6. Favus corporis, . . . . .	13
7. Trichophytosis capitis, . . . . .	14
8. Trichophytosis disseminata, . . . . .	15
9. Favus capitis, . . . . .	16
10. Contagious impetigo, . . . . .	22
11. Confluent lesions of impetigo, . . . . .	23
12. Impetigo—umbilicated lesion on forehead, . . . . .	24
13. Psoriasis—punctate lesions covered with white scales, . . . . .	28

FIG.	PAGE.
14. Psoriasis—guttate lesions with scales washed off, . . . . .	29
15. Nummular and circinate psoriasis, . . . . .	30
16. Psoriasis—patches healing in centre, enlarging and coalescing, . . . . .	31
17. Ichthyosis, . . . . .	36
18. Ichthyosis with broken epidermis, . . . . .	37
19. Ichthyosis with fine scales, . . . . .	38
20. Polygonal plates of ichthyosis, . . . . .	39
21. Eczema squamosum, . . . . .	42
22. Eczema squamosum, . . . . .	43
23. Eczema rubrum, . . . . .	44
24. Eczema impetiginosum, . . . . .	45
25. Eczema pustulosum, . . . . .	46
26. Linear papilloma, . . . . .	52
27. Linear papilloma, . . . . .	53
28. Nævus, . . . . .	56
29. Nævus, . . . . .	57
30. Nævus, . . . . .	58
31. Nævus of nose, . . . . .	62
32. Birthmark of unusual extent, . . . . .	63
33. Nævus of scalp, . . . . .	64
34. Nævus tuberosus, . . . . .	65
35. Nævus of vulva, . . . . .	66
36. Lupus vulgaris—a group of tubercles, . . . . .	67
37. Lupus vulgaris—tubercles forming a ring, . . . . .	68
38. Serpiginous lupus with central cicatrix, . . . . .	69
39. A favorite site of lupus, . . . . .	70
40. Lupus verrucosus (tuberculosis verrucosa cutis), . . . . .	71
41. Ulcerating lupus with atrophy of bone, . . . . .	72
42. Lupus erythematosus, . . . . .	73
43. Papular form of lichen ruber, . . . . .	75
44. Lichen ruber—papules on a favorite site, . . . . .	76
45. Plantar lichen ruber resembling eczema, . . . . .	76
46. Lichen ruber resembling ichthyosis, . . . . .	77
47. Lichen ruber resembling psoriasis, . . . . .	77
48. Rugous form of lichen ruber, . . . . .	78
49. Lichen ruber—chronic form of disease, . . . . .	79
50. Moniliform lichen ruber, . . . . .	80
51. Lichen planus, . . . . .	81
52. Lichen planus, . . . . .	81
53. Lichen planus, . . . . .	82
54. Erythema nodosum, . . . . .	85
55. Purpura, . . . . .	86
56. Molluscum, . . . . .	88
57. Keratosis follicularis, . . . . .	90
58. Keloid, . . . . .	91
59. Scabies, . . . . .	93
60. Pustules in scabies, . . . . .	94



PLATE I.



ALOPECIA AREATA.  
From the collection of photographs of  
Dr. GEORGE HENRY FOX.

## ALOPECIA AREATA.

---

ALOPECIA areata is an affection of frequent occurrence in childhood, and one which the family physician should be competent to recognize and to treat. To say that he should be able to treat a case as successfully as any specialist is not making a great claim, when we consider how little positive knowledge is possessed by the latter concerning the cause and cure of this not uncommon disease.

As the name indicates, alopecia areata implies the occurrence of baldness in spots; but all bald areas are not necessarily cases of alopecia areata. Before describing the affection itself it may be well to refer briefly to certain forms of baldness, areate or complete, which are often improperly classed as cases of the disease in question.

When infants are born without hair, and for years, perhaps, exhibit but a sparse capillary growth, coincident usually with imperfect dental development, this condition is not to be regarded as alopecia areata and the term is not apt to be applied.

When, as a result of injury, fright, or intense mental excitement, the hair falls suddenly from the scalp, or from the whole body in the case of adults, we have a condition which is by no means one of true alopecia areata, although in such cases the name is very apt to be misapplied.

When, from injury to a nerve, an irregular patch of baldness develops upon the portion of hairy skin which the nerve supplies, the condition is one which strongly resembles the disease in question. It differs, however, from true alopecia areata in several essential points. It usually lacks the rounded outlines, it does not tend to spread or duplicate itself at other points, and the hairs at the margin of the bald area are not loosened. It is a bald patch of evident neurotic origin, but should be distinguished from alopecia areata.

A rare form of scalp disease, known as folliculitis decalvans,

occurs in children as well as in adults, and is very apt to be mistaken for the commoner disease, to which it may bear a strong resemblance. Its distinguishing features are the signs of inflammation in incipient patches, the groups of dark plugged follicles, and the atrophied or cicatrized appearance of the bald patches, which are usually numerous and small. In very rare cases of ringworm of the scalp a smooth, bald patch results (Fig. 1). Whether this is to be regarded as a peculiar form of ringworm (so-called bald ringworm), or as a case of alopecia



FIG. 1.—Bald ringworm.

areata developing on the site of trichophytosis, is a vexed question.

In true alopecia areata we have as essential clinical features the development of the disease in one or more small, rounded patches; the extension of these patches by a loosening and falling of the hair at a portion or the whole of the periphery; and a marked tendency to spontaneous recovery. The first bald spot in this affection may occur upon any portion of the scalp, and is usually the size of a ten-cent piece or larger when first



discovered. It is unaccompanied by itching or any sign of inflammation, and may have existed for many weeks before cutting of the hair or a casual examination of the scalp brings it to light. In exceptional cases tenderness upon pressure, anæsthesia, hyperæmia, œdema, marked pallor or depression of the patch, are symptoms which have been noted, but ordinarily a partial or complete loss of hair is all that can be seen. The original patch sometimes remains single, but usually other patches develop near the original one, and frequently, by increase



FIG. 2. - Serpiginous form of alopecia areata.

in size and coalescence, denude a considerable portion of the scalp, as is shown in the accompanying plate and in Fig. 2. The irregular area from which the hair has fallen is smooth and soft, and, unlike a patch from which the hair has been epilated or shaven, the mouths of the hair follicles are scarcely perceptible. In some cases the bald patches are numerous, of varying size, and show little tendency to run together. The affection is by no means limited to the scalp, and in severe cases the eyebrows and lashes also suffer. In adults the bearded portion

of the face and other hairy parts may present bald areas or appear completely denuded.

When first discovered the patch is usually tending to increase slowly in circumference, and the hairs at the margin of the disc are so loosened in their follicles that several can be pulled out at a time without giving the patient the slightest pain. Often the loose hairs will only be found on one side of the patch, and extension in this direction is certain to follow. Examined beneath the microscope these loose hairs will be found to be atro-



FIG. 3.—Growth of white hair on patches.

phied toward the root, and in some cases they will break near the surface of the skin. This breaking of the hairs does not leave a short, stiff stubble like that which is characteristic of the patches of ringworm, but as the broken-off roots of the affected hairs in alopecia areata project from the scalp near the margin of the patch, they may be noted as short, fine hair tapering downward and sometimes suggesting a crop of exclamation points.

The extent to which alopecia areata will develop in a given case it is difficult to determine in advance, but sooner or later in



most cases, and especially in those cases occurring in childhood, the cause seems to have ceased to act and the hair returns. A fine, downy growth may be seen over a portion or the whole extent of the bald patch, and gradually the baldness disappears. In some cases, and especially in children with dark hair, the new growth is at first colorless and presents quite a contrast with the normal hair (Fig. 3). If this new hair persists in growing, it is certain to become pigmented in due time and assume a natural color. But unfortunately in some cases this new growth does not persist. It grows just enough to



FIG. 4.—Complete loss of hair.

delude one into a hope of speedy recovery, and then begins to loosen and fall out. Even a temporary growth, however, is encouraging and warrants the prediction that sooner or later the disease will come to an end. In many cases two or three months will find the hair growing naturally over the whole scalp, while in other cases the affection, through a stubborn persistence or a succession of relapses, may last a year or two. In adults a complete loss of hair is often irremediable, but in

children the prognosis is comparatively favorable even when the head is as hairless as a billiard ball (Fig. 4).

The etiology of alopecia areata is a vexed question. Despite the great amount of study and careful observation which has been devoted to this interesting affection, there is much difference of opinion as to its pathology, and its true nature still remains obscure. For a long time it was believed to be a neurotic affection, the bald patch resulting from a faulty innervation of the affected area. With the development of the germ theory of disease there has arisen a strong belief on the part of many dermatologists that the affection is parasitic in character. Clinical facts are not wanting in support of either view, but, while the facts are indisputable, it cannot be admitted that they prove beyond all shadow of a doubt the truth of either the neurotic or parasitic hypothesis. Recently it has been suggested, in view of the conflicting facts and theories, that there are two varieties of the affection, one of which, occurring in epidemic form, must certainly be parasitic. Without attempting to introduce evidence and argue the question, it is sufficient to state that the nature of this disease is still *sub judice* and will doubtless so remain until more careful observations have been made or stronger arguments presented upon one side or the other. The disease occurs in both sexes, but is said to be somewhat more frequent among boys than among girls. It has been claimed that cats and other domestic animals suffer from the affection, and cases have been reported of its transmission from these to their human playmates; but in this connection the question naturally arises as to the correctness of the diagnosis between alopecia areata and ringworm, which not infrequently originates in this manner.

In the treatment of alopecia areata no brilliant results have been obtained. Many cases recover spontaneously in a short time, while in other cases the most assiduous and persistent treatment fails to effect any notable result. Many remedies are recommended and certain methods of treatment are warranted to cure in so many weeks, but, however valuable these methods may be in the hands of their originators, they often fail utterly when employed by other hands. I have photographed many cases and have observed that a few of these have gotten well speedily without further treatment. But photography will not cure all cases; neither will some of the vaunted methods of treatment, as I have had occasion to learn from experience. In fact it must be frankly confessed that the

treatment of alopecia areata is in many cases by no means as satisfactory as one might wish. Patients sometimes go from one physician to another, and fortunate is he who happens to take charge of the case at the time when the unknown cause has ceased to act and the hair is disposed to return. Unfortunate is he, however, if he jumps at the baseless conclusion that his plan of treatment in this case is wonderfully efficacious and will cure other cases with equal rapidity.

But treatment in alopecia areata, although it may not be productive of brilliant results, is capable of doing some good in the way of hastening a cure ; and since the patient usually demands some kind of treatment, the physician is certainly justified in doing the best he can under the circumstances. The practical question at once arises, "What is the best thing that he can do?"

The first step to be taken is to prevent any possible spread of the disease to those who come in contact with the patient. If only in deference to the view held by excellent authorities that the affection is of a contagious nature, it is well to insist that every patient should have his or her own brush and comb and use no others. As to whether children with alopecia areata should be allowed to attend school, a difference of opinion might arise according to the opposite views held as to its contagious nature. But since the disease is only slightly, if at all, contagious, it would hardly seem justifiable to keep a child at home who is undergoing proper local treatment.

Whatever will stimulate the scalp and draw the blood to the pale, hairless patches will tend, in my opinion, to promote the growth of hair. And this can be done without the infliction upon the patient of either pain or discomfort. Nearly all of the powerfully stimulating applications are parasiticide in their action, and are therefore indicated, whether one believes in the neurotic or the parasitic origin of the disease. The beneficial effect results from the artificial congestion of the patch, whether the action of this be to arouse the dormant nerve filaments or to check the development of a micro-organism.

The galvanic current is perhaps the best stimulant that can be used ; but as the patient cannot conveniently have electricity applied two or three times a day, it is more advisable to prescribe some stimulating lotion or ointment. The liquor ammoniæ fortior U. S. P. I have used for many years and in many cases ; and, while it may be no better than other stimulating applications which tend to redden the skin, it has certainly



appeared to do as much good as any one of the many other remedies which I have tried. If something more parasiticial in its action is desired, an ointment of chrysarobin from three to ten per cent in strength can be recommended as an effective application. In prescribing this the physician must not forget to mention the fact that it will stain the bed linen, and caution the patient not to get any ointment in his eyes lest a severe conjunctivitis result.

The plan long recommended and often practised of shaving the scalp and blistering the patches with cantharis, croton oil, or pure carbolic acid, may be of some value, but not enough to compensate for the suffering which it involves. Although patients will sometimes stand any amount of painful treatment in order to get cured, there is no justice in submitting them to harsh measures when milder ones are of equal value.

Epilation of the loose hair at the margin of the bald patch is advisable, since it seems in some cases to prevent the spread of the disease; but general epilation is unnecessary, and, so far as I know, is never practised in America.

A patient seen recently by the writer has had alopecia areata for ten years. Two months after marriage his wife found three bald spots upon her own head. At a European health resort the physician who was consulted, evidently believing in the parasitic origin of the disease, ordered the hair to be epilated not only on the scalp but on other hairy parts of the body. The spots persisted, and complete epilation was repeated in the case of both husband and wife. The lady's maid complained of slight falling of the hair, and her scalp was ordered to be thoroughly epilated, in spite of her indignant protest. In these cases, which illustrate the absurd extreme to which theoretical therapeutics may be carried, the vigorous treatment would have been pardonable had it been successful, but the gentleman, an extremely nervous individual, has still bald patches upon his scalp and chin.

Of internal medication it need only be said that neither arsenic, jaborandi, nor any other drug in the pharmacopœia is capable of producing any manifest result in most cases. They may be of some value, as has been claimed, but reliance upon them is usually disappointing. The restoration of hair after administration of a drug is no proof of its value in a single case. The absence of improvement after its use in repeated cases is at least suggestive of its inertness. Any internal medication which will improve the physical condition of the patient is of

course advisable and may tend to hasten the cure of alopecia areata. But in this affection impairment of health rarely seems to be a predisposing cause, and in many cases the patient is apparently in perfect health.

The uncertain result of treatment in alopecia areata should always lead to a guarded prognosis. In most cases a cure can be promised, but it is never advisable to specify any given time at which it may be expected. The younger the patient, the less the extent of baldness, and the more recent its development, the better will be the prospect of a speedy return of hair.





PLATE II.



TRICHOPHYTOSIS CAPITIS.  
From the collection of photographs of  
Dr. GEORGE HENRY FOX.



## RINGWORM AND FAVUS.

---

THERE are few if any cutaneous affections so ineffectually and unsuccessfully treated by the physician in general practice as are ringworm and favus. The family physician who assumes to understand and undertakes to treat (as every family physician should do) the common affections of the skin is often guilty of serious malpractice in the management of these parasitic affections. The diagnosis is generally made without difficulty, although mistakes will sometimes occur in the best-regulated practice. But, granting that the diagnosis is speedily and correctly made, the treatment commonly employed is so poorly adapted to the requirements of the case and so imperfectly carried on, that many incipient cases, which ought to be cured by a few weeks of vigorous treatment, are allowed to persist for months or years, it may be, and to become so chronic and intractable that even the most skilful treatment must then require a long period of time and much patience to effect a cure. Nor is this the only baneful effect of this feebleness of therapeutic efforts. Other children in the family, the school, or the vicinage are permitted to contract the disease, while the physician treating the original case remains blissfully ignorant of the ever-increasing debt of suffering which is his just due. Therapeutic incompetence may appear less criminal than wilful neglect, but the results are equally unfortunate, and in the treatment of these common parasitic affections the physician ought not to be guilty of either charge.

There are three common affections of the skin due to the presence of a vegetable parasite—viz., favus, trichophytosis (ringworm), and chromophytosis (tinea versicolor). As the latter is rarely if ever met with in childhood, it may be left out of consideration at present; and since the remaining two, though distinct in origin and clinical features, are allied in nature and call for similar methods of treatment, it will be convenient to consider them together.

Upon non-hairy parts, such as the face, neck, hands, and trunk, ringworm begins as a minute, slightly reddened, scaly disc, which gradually enlarges. When it has reached the size of a ten-cent piece the advancing border appears elevated while the centre tends to become smoother, and the lesion presents an oval, circinate, or "ring"-like appearance (Fig. 5). In rare cases a circle of fine vesicles can be detected at the periphery. One or more rings may be present, and the coalescence of two



FIG. 5.—*Trichophytosis corporis*.

or more may produce an irregular patch like a figure of eight or a trefoil. Occasionally when the ring has attained considerable size red papules or new foci of disease may appear in the smooth and perhaps slightly pigmented centre, and by gradual development may produce two or even more concentric rings.

Favus developing upon non-hairy parts produces scaly discs which at first are not readily distinguishable from the lesions of ringworm; but soon upon the branny surface a minute yel-

low, cup-shaped crust of the size of a pin's head will develop, which at once settles the question of diagnosis. These bright-yellow, pin-head cups multiply and by coalescence form a sulphur-colored crust which is quite characteristic (Fig. 6).

Ringworm and favus of non-hairy parts cannot be considered as serious affections, but when the parasitic fungus of either finds its way into the hair follicles upon the scalp, as often happens with children, the case is quite different. The unfor-



FIG. 6.—Favus corporis.

tunate child is now the victim of an extremely obstinate disease, and, unless judicious measures are adopted without delay, is doomed to months or perhaps years of annoyance, if not of actual suffering. Upon the scalp these affections are always obstinate, and when this fact is not appreciated by the physician in charge of the case, and inefficient measures are adopted, the case usually goes from bad to worse until a cure seems nearly hopeless.

The first indication of ringworm of the scalp is commonly a

small scaly disc, which appears almost bald from the breaking of the hairs close to the surface (Fig. 7). This dry, roughened patch, with its characteristic growth of short, broken hairs, tends to enlarge if allowed to go untreated (see plate), and other discs are apt to develop in the vicinity or upon other portions of the scalp. Frequently a large number of scaly points or small discs may be found involving the greater portion of the scalp and constituting what is known as disseminated ring-



FIG. 7.—*Trichophytosis capitis*.

worm (Fig. 8). Neglect in such a case is usually followed by a coalescence of the patches and disease of nearly the whole scalp. Not infrequently an eczema complicates the ringworm and obscures the diagnosis.

In rare cases the ringworm fungus, instead of producing scaly discs, sets up a deep-seated inflammation of the hair follicles, and a bald, fluctuating tumor or a cluster of boggy, rounded elevations forms upon the surface of the scalp. This condition is generally painful, and is known as kerion or the



kerionic form of ringworm. The suppuration often loosens the hair, and in these cases complete baldness is more frequently found than the stubble-like growth of hair which characterizes the ordinary form of the disease. According to recent observations, this form of ringworm is due to a special fungus which also affects the lower animals.

In favus of the scalp there are no broken hairs found as upon the discs of ringworm, and the diagnosis is usually based upon



FIG. 8.—*Trichophytosis disseminata*.

the presence of the minute yellow, cup-shaped crusts which develop at the orifices of the hair follicles. When these are allowed to multiply, a thick, pale-yellow, friable crust forms, as is seen in the well-marked case (Fig. 9) which was originally published in the author's "Photographic Illustrations of Skin Diseases." Much has been said about the peculiar odor of favus as a basis of diagnosis. While it is true that in a neglected case, where the crusts have accumulated, a keen scent combined with a vivid imagination may perceive a fragrance which is

quite suggestive of "mice," "putrid urine," or "an ill-kept menagerie," the diagnosis can be far more readily made from the characteristic appearances. In doubtful cases, moreover, in which there are no characteristic crusts, the peculiar faveic odor is not likely to be perceived by ordinary olfactories. While typical cases of ringworm and favus are totally unlike in their clinical appearance, treatment often obscures the characteristic features, and hence, in cases where the crusts have



FIG. 9.—Favus capitis.

been removed, the differential diagnosis may sometimes be attended with difficulty. In ringworm, however extensive and chronic the case may be, the hair is not apt to be permanently destroyed; while in favus the pressure of the crusts which develop in the epidermic layer around the orifices of the follicles tends to produce atrophy of the hair bulbs, and in all cases of long standing a few bald, depressed, and cicatricial areas are generally observed, and upon these the hair will never grow. In chronic cases, even after the disease has been cured, a num-

ber of wiry, twisted, or deformed hairs are usually seen growing around or among these cicatricial patches.

Trichophytosis capitis is a disease of youth, and even when allowed to go untreated it will tend to a spontaneous cure as the patient matures. Though ringworm of the beard is very common in men, for some inexplicable reason ringworm of the scalp is never met with in adult life in either male or female. Favus, on the other hand, occurs at all ages. Developing in youth, it may persist indefinitely, although it is far more frequent in childhood than in adult life.

The cause of ringworm and favus is the presence in the epidermis and hair follicles of a micro-organism, the growth of which gives rise to more or less inflammation and the characteristic clinical appearances already described. The parasitic fungus causing ringworm is the *trichophyton*. Several varieties have recently been described, but it is not as yet satisfactorily proven that these varieties of the fungus are the cause of all clinical variations noted in the course of the disease. The etiological factor in favus is the *achorion*. This fungus does not penetrate the shaft of the hair as readily as does the trichophyton, and hence the absence in favus of the broken hairs which are so characteristic of ringworm. Children in perfect health are liable to contract either disease from some child already affected, or possibly from some pet animal. No particular condition of the skin is necessary to furnish a congenial soil for the development of the parasite, although it is true that in weak, poorly nourished children whose heads are often moist the disease is more likely to thrive and the inflammatory symptoms to be more marked. Age, however, seems to modify the character of the soil in a notable degree, and it is well to remember the clinical fact, already mentioned, that ringworm of the scalp, so common in childhood, is never seen in adult life. In like manner ringworm of the beard, so frequently seen in middle life, is quite exceptional among old men.

In the treatment of ringworm and favus a host of local applications are recommended by dermatological writers, which only proves that the majority of them are of very little value. Moreover, an admirable prescription may be written, but if the nurse or person in charge of the patient is not given full and explicit directions as to how the treatment should be carried out, the result is certain to be unsatisfactory. The local remedy employed is often of far less importance than the exact method of its use, and attention to minute details which may seem



unimportant to the inexperienced is always the key to success. In no other affections of the skin are intelligence and persistence so essential, and for the exercise of these in the daily care of the patient the physician should hold himself responsible.

On non-hairy parts ringworm and favus can be readily cured by almost any parasiticide. A ten per cent ointment of salicylic acid is an effective application, or, if the skin is very delicate, it may be better to simply moisten the patches frequently with a saturated solution of sodium hyposulphite in rosewater. But when the scalp is affected the cure is always a difficult one, and the first step is to impress upon whoever is in charge of the case the important fact that half-way measures will do little or no good. It is always advisable to shampoo the scalp thoroughly once a day, especially if the hair is short, as this gives the parasiticide application a much better chance to penetrate the hair follicles. The neglect of frequent and thorough washing of the scalp is the chief cause of the frequent therapeutic failures. When this is carefully attended to, the thorough inunction twice daily of sulphur ointment, oleate of mercury, or chrysarobin ointment (ten per cent) is certain to do good. If a cap is worn by the patient, as is advisable, it should be fastened to the head by a ribbon or strip of bandage, and not by a rubber band, on account of the serious results which might ensue from a prolonged stoppage of the circulation.

There is one remedy which is indispensable in chronic cases and of the greatest value in any case. It is epilation. It must be admitted that this is always troublesome to the physician or nurse who undertakes to carry it out, and more or less painful to the patient. But it saves time and trouble in the end. With a well-made pair of epilating forceps, which should be light and broad at the end of the blades, the short hairs can be firmly caught and quickly pulled out. It is advisable to epilate first the long hairs around the margin of a patch until a narrow white ring of healthy scalp appears. This will prevent any increase in size of the patch, and the short hairs upon the reddened, scaly surface of the patches can be pulled at leisure. As many of these will break in the process of epilation, the operation must be repeated until the patch is quite bald and begins to assume a comparatively healthy appearance. The epilation and the application of parasiticides can be carried on at the same time. When the inflammation has subsided and the scaling disappeared, and all the patches have assumed a comparatively healthy appearance, the hair may be allowed to



grow and all treatment suspended save the daily use of a five per cent salicylated oil. If, however, at any time a slight scaliness or dry, brittle appearance of the hair is noted at any point, it is advisable to epilate again and convert the suspicious spot into a small bald disc. This plan of treatment is best calculated to effect a certain if not a speedy cure, but often it will require months of patient and persistent treatment, and perhaps a year or more in exceptionally extensive and chronic cases.

The parents or guardians of the patient should always be apprised at the outset of the obstinacy of the disease and its unfavorable prognosis as regards a speedy and pleasant cure, in order to avert the discouragement and dissatisfaction with the method of treatment which otherwise would naturally ensue.

It seems hardly necessary to add that no child with ringworm should be allowed to attend school. If our city Board of Health could make provision for a periodical inspection of the public-school children, with a view to checking the spread of ringworm and other contagious diseases, a considerable amount of suffering and expense could be saved to a certain number of scholars and their parents. Furthermore, if some careless physician were sued for malpractice for allowing an uncured case of ringworm or favus to attend school, simply because upon a hasty inspection, without the use of a microscope, he thought the child was all right, it might be unfortunate for the physician, but by no means a bad thing for the profession.





Plate III



N.Y. PHOTOGRAPHIC CO.

*Impetigo Contagiosa*

From the collection of photographs of

*Dr. George Henry Fox*

## CONTAGIOUS IMPETIGO.

---

THE affection known as contagious impetigo, though a very common one, is by no means as clearly defined as other dermatoses of frequent occurrence. Crusted lesions or scabs resulting from the use of the finger nails and the drying of a copious blood-stained secretion are frequently seen upon the faces of children, but the origin and nature of this eruption are not always plain. That the lesions are contagious and auto-inoculable is a matter of simple observation, and that pus cocci play an important part in their development is generally admitted. Mothers and nurses are often affected by children under their care.

It is undoubtedly the presence of micro-organisms which occasions the crusted or impetiginous form of eczema (the crusta lactea, or milk crust of infants). In these cases we have an ordinary eczematous inflammation of the skin due to some internal condition, but the peculiar clinical aspect of the eruption—viz., the suppuration and crusting—is without doubt due to the external agency of pyogenic microbes. In other cases minute aggregated pustules develop superficially and exude an abundant honey-like secretion which dries into yellow, gummy crusts of varying size and irregular outline. This eruption is comparatively rare, affects adults as well as children, and is by some regarded as a pustular eczema and by others as impetigo or contagious impetigo. In fact it is difficult to determine in practice precisely what cases should be included under the last-named affection, but a large number of cases occurring in children present certain definite lesions which are readily recognized and indicate an identity of origin.

In a typical case of contagious impetigo (and it is questionable whether there is any impetigo which is not contagious) the lesion is usually a flattened vesico-pustule. This is small at the outset, superficial, and without any surrounding area of inflammation. It is always single, although when the pustules

are very numerous, a number of them in close proximity may appear like an irregular group. When this lesion escapes being scratched, and thereby altered in appearance, it tends to increase peripherally and to become depressed in the centre. It usually reaches its maximum development in a few days, and is about the size of a five-cent nickel. At the advancing border the epidermis is raised by a milky, purulent secretion while the depressed centre appears of a darker hue. Upon the back of the fingers or hand, where, next to the face and scalp,



FIG. 10.—Contagious impetigo.

the lesions are most common, they look like recent and accidental burns. Upon the palm, where the thicker epidermis is less likely to be ruptured, the lesions are apt to present the appearance of hemispherical or flattened pustules, quite similar to, though somewhat larger than, the pustules occasionally observed in scabies. After an existence of about a week the contents of the pustules tend to dry into yellowish, honey-like or "straw-colored" crusts. If the lesions remain uninjured by the finger nails, the affected skin gradually heals and the crusts



appear as though "stuck on" to the cutaneous surface. When thoroughly dried these crusts fall, and even when picked or rubbed off they leave no ulcerated surface, but merely a small hyperæmic area.

It is only in exceptional cases, however, that this natural and characteristic evolution of the lesion can be observed, for in children, especially, the irritation of the lesions usually provokes scratching, which changes their clinical appearance, and a little blood mingled with the purulent secretion gives rise to



FIG. 11 —Confluent lesions of impetigo.

the formation of a dark crust or scab. Upon the extremities, especially the legs, friction of the clothing, dirt, and constant scratching frequently inflame the lesion, and a dark crust upon a superficial ulceration with a narrow red areola is produced, which has generally been described as a distinct skin affection under the name of ecthyma. These ecthymatous pustules, common in cachectic children, are often associated with the characteristic lesions of contagious impetigo upon the face and hands and are evidently of similar if not identical origin.

In certain cases of vaccination a double infection may sometimes take place. When the vaccine virus has produced a vesicle, the microbes of contagious impetigo may be accidentally implanted. The vaccine lesion now becomes unusually inflamed and itchy. Through scratching a number of other crusted lesions are developed upon the arm and elsewhere, and doubtless some of the cases of generalized vaccinia which have been reported may be justly considered to have been cases of contagious impetigo beginning at the point of vaccination.



FIG. 12.—Umbilicated lesion on forehead.

Upon the scalp the lesions of contagious impetigo are frequently found, and usually appear as small, isolated, circular crusts. These frequently occasion temporary areas of partial baldness, and when the disease is of long standing and there is very much suppuration a permanent loss of some hair may result. The affection is often associated with pediculosis capitis, and in such a case the frequent scratching of the head is very apt to multiply the lesions, and often the diagnosis is obscured by the development of eczema upon the scalp and back of the



neck. In impetigo of the head and face the cervical and sub-maxillary glands are very apt to become enlarged and tender, and in strumous, ill-nourished, and neglected children glandular suppuration may occur. Among this class the affection frequently coexists with paronychia and ulcers of the conjunctiva and buccal mucous membrane. It is said to occur with especial frequency in female children and in the spring, although neither sex nor season is exempt.

The upper illustration in the accompanying plate shows crusted lesions upon the forehead and scalp, and a secondary conjunctivitis which has closed the right eye. Upon the hand a number of incipient pustules are seen, excoriated upon the fingers, but tolerably well developed upon the wrist. Upon the forearm, in the lower illustration, a few typical flattened and umbilicated lesions may be seen, while upon the arm an ecthymatous patch has resulted from the coalescence of several scratched and crusted lesions. In Fig. 10 the eruption presents its characteristic appearance in the form of numerous isolated crusted lesions. In Fig. 11 a few lesions have coalesced upon the lips and formed a large, crusted patch resembling impetiginous eczema. In Fig. 12 the lower lip is similarly affected, while upon the forehead the isolated lesions present a typical appearance.

While the individual lesions of contagious impetigo run an acute course of a week or ten days, the development of new lesions and the irritation produced by scratching often keep up the eruption for many weeks. In its treatment cleanliness is of the utmost importance, especially as the affection is most likely to occur among children whose ablutions are neither frequent nor thorough. The finger nails may be advantageously cleansed and cut, and a disinfectant lotion, such as listerine or a saturated solution of sodium hyposulphite in rosewater, should be applied frequently to the affected skin. To remove crusts and heal superficial ulceration an excellent application is a mixture of equal parts of white precipitate ointment and cold cream.







*Psoriasis*

*From the collection of photographs of*

*Dr. George Henry Fox.*

## PSORIASIS.

---

PSORIASIS, though most frequently met with after the age of puberty, is by no means a rare affection in childhood and may even occur in infancy. Though different cases present a notable variation in their clinical appearance, the characteristic features of the disease are very much the same at all ages. The eruption is always dry and scaly, whatever may be the age of the patient, but in childhood it is not generally so well developed and so extensive as it is apt to be in later years.

Psoriasis begins in the form of one or more red points, which quickly become covered with white, silvery scales (Fig. 13). These may be readily scratched off by the finger nail, and when this is done a bleeding surface is exposed. When many of these small, scaly lesions are present the eruption is described as punctate psoriasis, and this form of the eruption is comparatively more frequent in children than in adults; when the scaly lesions increase in size and appear like drops of grease or thin mortar spattered over the skin, we have the guttate form of the disease (Fig. 14); and when the patches assume the size and shape of silver coins they are often described as nummular psoriasis (Fig. 15). By healing in the centre these lesions may be converted into scaly rings, or by peripheral increase and coalescence they may result in the formation of extensive scaly patches (Fig. 16). Diffused or general psoriasis is, however, rarely met with among children.

The amount of scaliness present in any case depends upon the attention which the patient naturally devotes to his skin. If baths are frequently taken, and especially if any fat or oil is rubbed over the patches, the scales are generally absent and the eruption presents a tolerably smooth, reddened appearance. When no attention is paid to the care of the skin, the scales often accumulate upon the psoriatic patches until they are very prominent and present a silvery-white or dirty-yellow appear-



ance. As the eruption tends to disappear the scaling grows less, often disappearing from the centre of the patch and leaving a marginate ring. Finally the redness fades and the skin assumes a normal appearance, except in certain cases where pigmentation may occur.

In rare cases of psoriasis the eruption may tend to rapidly involve the whole skin. The cutaneous congestion is severe, and large flakes of partly detached epidermis may take the place of the silvery scales. In other words, an attack of acute



FIG. 13.—Punctate lesions covered with white scales.

dermatitis exfoliativa has set in and complicated and obscured the psoriasis. When the dermatitis has subsided the psoriasis is very apt to reappear.

Psoriasis is not only a dry and scaly eruption, but it is invariably marginate. Whether occurring in small discs or in large, irregular patches, the border is always sharply defined and never shades off gradually into the surrounding healthy skin, as does the ordinary patch of eczema. This is a diagnostic point of great importance. In many cases of eczema the

patches may be dry and scaly, and present a resemblance to those of psoriasis, but the rounded, silvery discs or larger marginate patches of the latter disease are usually so characteristic that an error in diagnosis is not likely to be made.

The localization of the eruption is another important diagnostic point. While eczema may appear upon almost any part of



FIG. 14.—Guttate lesions with scales washed off.

the body, and often exhibits a tendency to attack the flexor aspect of the joints and other parts where the skin is thin and delicate, psoriasis is commonly seen upon the extensor surface of the extremities and is especially apt to be noted about the elbows and knees. Upon the scalp the two affections often present a strong resemblance, but in psoriasis the scaly patches



are apt to be small, numerous, and circular, with healthy skin intervening, while eczema of the scalp usually occurs in one large, diffused patch.

The symmetry of the eruption in psoriasis is also a characteristic feature. The eruption upon one extremity or one side



FIG. 15.—Nummular and circinate psoriasis.

of the trunk is usually duplicated upon the other side, while eczema is very frequently unilateral.

The papular syphilide often assumes a squamous form and presents for a time a strong resemblance to psoriasis. Fortunately it is rarely met with in childhood, but at any age it

differs from psoriasis in one respect. While in the latter affection the infiltration of the skin is slight and the accumulation of scales a prominent feature, in the papular syphilide the infiltration of the skin is considerable and the scaling is comparatively slight.

It is a noteworthy fact that many patients suffering from



FIG. 16.—Patches healing in centre, enlarging and coalescing.

psoriasis are unusually well developed and robust in appearance. It is not, like eczema and certain other skin affections, a disease of the weak and the neglected, but one which seems to delight in attacking those who are strong and well nourished. In any case, however, where the tendency to psoriasis exists, the eruption is certain to be most severe whenever the patient is weakened by lack of proper food or by overwork.

The tendency to psoriasis is frequently inherited, and often the disease may be observed in two or more generations. Not infrequently it shows a tendency to skip a generation, and it has been claimed that psoriatic subjects are the offspring of eczematous, dyspeptic, asthmatic, gouty and rheumatic, as well as of psoriatic parents. External irritation only produces the disease in those who manifest a predisposition to it.

While it is often an easy matter to remove the eruption by treatment, it is difficult, if not impossible, in many cases, to prevent its speedy return. The eruption shows a marked tendency to relapse or to increase in severity in the spring or at some particular season, and in many cases persists for years or even throughout a lifetime. It often improves or disappears spontaneously for a few months or even for a year or more, and then reappears and continues its fluctuating course. The prognosis, therefore, is always a grave one as regards the permanent cure of the disease; but, in spite of the fact that some have declared psoriasis to be incurable, there are many cases which get well and remain well.

In the treatment of psoriasis a host of remedies have been employed, and many of them owe their repute to the fact, already stated, that the eruption tends at times to disappear spontaneously, and to any remedy employed at such a time the credit of a cure would naturally be awarded.

Among internal remedies, arsenic is the one most commonly employed and probably the most efficacious. In many cases, however, it may do much more harm than good. When the skin is irritable and the psoriatic patches congested, arsenic is worse than useless and alkaline diuretics are greatly to be preferred. But, on the other hand, when the disease is tending to get well, the administration of arsenic will often produce a most brilliant therapeutic result.

Of the various local remedies employed, chrysarobin stands without a rival. In many cases it produces a speedy and brilliant result which can be attained by no other local application. It has serious objections, however, which often forbid its use. When the skin is irritable it may cause the eruption to spread. Like arsenic, it is most likely to do good when the acute congestion of the psoriatic patches has subsided and the eruption is tending toward a spontaneous improvement. It not only stains the skin temporarily—which, however, is a matter of little importance—but it permanently discolors the underclothing and the bed linen, if due precaution is not taken. When

rubbed in where the skin is thin, or near it, as, for example, the axillæ and flexures of the joints, it often excites a very unpleasant dermatitis for a few days; and when by chance a little of the ointment gets into the eye a very severe conjunctivitis often results. This chance is somewhat lessened by applying the drug in the form of a varnish composed of gutta-percha solution or collodion, but then its therapeutic effect is decidedly lessened. Upon the trunk and extremities a five or ten per cent ointment, made by rubbing up *finely sifted* chrysarobin in vaseline, can be advantageously used; but upon the scalp and face the ointment of ammoniated mercury will generally prove efficacious, and is to be preferred to the chrysarobin.

Prolonged baths are often of service in macerating the scaly patches, and, when the skin is not too irritable, soap frictions are valuable in removing the scales and preparing the skin for inunction. In many cases, however, owing to the intense congestion of the affected skin, a hot bran or alkaline bath will prove more agreeable than the use of soap.










Plate V.



N.Y. PHOTOGRAPHIC CO.

*Ichthyosis*

*From the collection of photographs of*

*Dr. George Henry Fox.*

## ICHTHYOSIS.

---

ICHTHYOSIS is a deformity or an imperfect development rather than a disease of the skin. Like all deformities, it is very apt to be hereditary, affecting one or more members of a family, and often skipping a generation. It is characterized by a marked deficiency of the normal cutaneous secretions and a tendency to the formation of a dry, scaly surface, which suggested the name of "fish-skin disease." It is sometimes congenital, may indeed develop *in utero*, but often does not manifest itself until several months after birth. In rare instances it may develop late in life. Though always improved by judicious treatment, it may be considered as one of the incurable dermatoses, since the peculiar character of the skin cannot be changed.

There are various degrees of severity in which ichthyosis may manifest itself, and different names applied to these have led to some confusion as to their true character. For instance, the term xeroderma, or "parchment skin," has been used unnecessarily to designate a very mild form of ichthyosis in which the skin presents a dry, mealy appearance, especially apt to be noticed in children with whom bathing is a rare luxury. In some cases this slight ichthyotic tendency disappears in time, and sometimes appears in later life as an acquired condition. It should be borne in mind that as the hair in some children is preternaturally dry and in others unusually oily, and that as this condition may vary in the same individual according to the state of health, so the degree of natural oiliness or dryness of the skin may vary greatly in individuals and in accordance with season and general condition. A skin which is notably dry and with a tendency to chap or roughen might be regarded as xerodermatous or ichthyotic in a very slight degree.

In ichthyosis mitis (or simplex) the characteristic features of the affection are well developed. The skin is dry, and perspi-

ration is slight even in warm weather. Upon the extensor aspect of the extremities the epidermis presents often a peculiar serpentine appearance, resulting from the cracking of the horny layer into polygonal plates, which in time assume a dirty gray or greenish tint. These plates adhere by the central portion,



FIG. 17.—Ichthyosis.

while the margin tends to separate from the underlying skin. Around the elbows and knees the natural wrinkles are greatly intensified, and the breaking of the horny epidermis occasions a large number of concentric, whitish lines separating rows of warty elevations. As a rule the flexures of the joints, axillæ, groins, genitals, palms, and soles are but slightly if at all



affected. The face usually presents a characteristic appearance, the cheeks being roughened or chapped, the eyelids stiffened and drawn into a condition bordering on ectropion, the lips dry, and the oral commissures more or less wrinkled. The hair is usually dry and sparse upon the scalp, and may be entirely absent upon those portions of the trunk and extremities where a slight growth is usually present. The nails are apt to be dry and brittle.

In cases of ichthyosis of a severe type in which no treatment is instituted and the ordinary use of soap and water is neglected, the epidermis becomes broken into small plates, which increase in thickness until the affected skin upon the extensor aspect of




FIG. 18 — Ichthyosis with broken epidermis.

the extremities is covered with more or less conical, blackish masses and resembles the bark of a tree. To this condition the term ichthyosis hystrix, or "porcupine skin," may be applied. It should be borne in mind, however, that this term is often used to designate a linear warty condition occurring in parallel streaks, usually upon one side of the body (*nævus verrucosus*), and which has no clinical relation to ordinary ichthyosis.

The severest form of the disease is one which develops *in utero* and is frequently fatal shortly after birth. This has been described as "diffuse congenital keratoma," "harlequin foetus," etc. Dr. George G. Wheelock in a report of a case<sup>1</sup>

<sup>1</sup> Illustrated Quarterly of Medicine and Surgery, July, 1882.

states that the forceps were applied through the os and the child was with considerable difficulty extracted, there seeming to be a total lack of lubrication and consequently great friction between the foetal and maternal parts. As the head was born a thick plate of skin two inches square was detached and escaped with the head. At first the child had the appearance of a dead foetus with macerated epidermis, but shortly it began to breathe and to cry feebly. Its appearance was horrible in the extreme. It was covered from head to foot with a skin like leather, deeply fissured and broken up into plates like an alligator or an armadillo. Many of the plates were separated from the true skin, which was of a bright strawberry color. After birth the dried



FIG. 19.—Ichthyosis with fine scales.

skin became of a bright chrome yellow, and the plates were more and more detached by the motions of the child, which lived only six hours. Another case has been reported of a woman who had three healthy children by her first husband and three ichthyotic foetuses by her second. Cases have also been reported of this severe form of the disease developing after birth.

Occasionally one of these cases of severe congenital ichthyosis may survive. A youth known to fame as the "Alligator Boy," and exhibited in a dime museum some years ago, was described by the writer in the *Journal of Cutaneous and Venereal Diseases*, April, 1884. The eruption, contrary to the rule, was most marked upon the trunk, the epidermis being

broken by movements of the body into polygonal, horny plates of varying size and of a dirty-yellowish hue. In winter the skin upon the trunk was smooth, though thickened and horny in character. In the spring this horny integument was wont to crack and present the characteristic alligator appearance,




FIG 20. —Polygonal plates of ichthyosis.

while the hair of the head was almost entirely shed. At birth, according to Dr. F. Kennedy, the child presented a most extraordinary appearance. The skin was smooth, as if polished, and of a deep-red color. It was impossible for the child to make use of its facial muscles in its efforts to cry or suck. After a



few days fissures occurred and armor-like scales were formed, which gradually peeled off. This case was evidently of the same nature as those reported by Wheelock, Kyber, and others, but not so severe as to cause death.

Heredity is the only cause which can be suggested for the occurrence of ichthyosis, but often the parents and grandparents are said to have been free from this or any other form of skin disease. The sexes are about equally disposed to the affection.

The treatment of ichthyosis is of necessity palliative, as the cause cannot be removed, but even in the severest cases the skin can be softened and restored temporarily to an almost or quite normal condition. To maintain this condition a considerable amount of persistent daily lubrication of the skin is required. The use of fatty articles of food, such as cream, eggs, etc., is to be recommended; and cod-liver oil will often prove serviceable, although it may not be necessary to use it internally, externally, and eternally, as has been suggested. Indeed, for local application the fats and oils do not generally prove as agreeable and effective as does diluted glycerin. A half-ounce of this added to a pint of rosewater (or rainwater) makes an excellent lotion, which can be readily rubbed over the extremities, or whole body if necessary, every night and morning. When the disease is complicated by eczema, as often happens, a soothing ointment or paste is called for.





*Eczema.*

*From the collection of photographs of*

*Dr. George Henry Fox.*

## ECZEMA.

---

ECZEMA is at all ages the most important skin disease which the physician is called upon to treat. In infancy and childhood it is especially common, and usually a source of great discomfort to the patient. A full discussion of the subject would require much time and space, and in the few pages allotted to it here only a few points can be briefly mentioned and a few practical suggestions offered.

Eczema is an inflammation of the skin, of which the symptoms are commonly redness, thickening, moisture, crusting, and itching. The disease may present a great variety of appearances, according to the locality involved and the duration and severity of the inflammation. It may bear a resemblance to nearly every other skin affection, but in most cases, especially in childhood, the disease is readily recognized. Unlike many other cutaneous affections, it has no characteristic lesion, but may be erythematous, papular, vesicular, or pustular at the outset and gradually become crusted, scaly, fissured, or ulcerated. As regards the grade of inflammation, the disease may be observed in an acute, subacute, or chronic form, and as regards duration it may in one case be insignificant and ephemeral while in another case it may persist or recur year after year.

A multitude of adjectives have been applied to the numerous and varied clinical forms of eczema, and are useful for purposes of description, but the main thing for the physician to learn is to recognize the eczematous nature of the eruption, and to remember that different clinical forms may coexist upon different portions of the body and that one form is very likely to develop into another as the eruption becomes better or worse.

Eczema in childhood, as in later years, may occur in either an acute or chronic form. These terms, as commonly used, indicate the grade of inflammation rather than the length of time which the eruption has existed. In the acute form there is much redness, heat, and swelling, while in the chronic form



there is usually more thickening of the skin. An acute eczema often runs a typical course, which may be divided into three stages. These are, first, the stage of congestion and vesiculation; second, the stage of moisture and crusting; and third, the stage of desquamation. The vesicular condition is always transitory, and in many cases is not present. The effusion of serum may be so intense as to loosen and wash away the epi-




FIG. 21.—Eczema squamosum.

dermis in a mass, thus leaving the typical raw, red, exuding surface; or, on the other hand, the serous effusion may be so slight as not to appear upon the surface, and the eruption passes directly from the congestive into the squamous stage. The outbreak of acute eczema is attended by more or less fever and restlessness. Repeated exacerbations often characterize the course of a chronic eczema, during which the eruption assumes an acute form.



There are four types of eczema which it is well to differentiate and of which a brief description may be given. The first is erythematous eczema, in which redness and slight scaling are the chief features. It is usually a dry and mild form of the disease, and was formerly described as a distinct affection under the name of pityriasis. It is frequently seen upon the face and hands of children, and appears like a roughened or chapped con-



FIG. 22.—Eczema squamosum.

dition of the skin. About the genitals and wherever folds of skin are in apposition it is apt to follow the affection known as erythema intertrigo, and in many cases, indeed, it is difficult to distinguish between the two, as one gradually develops from the other. In erythema we have simply congestion of the skin, while in eczema erythematosum there is a slight thickening from serous infiltration of the cutaneous tissues and a tendency

to the development of a moist surface when the skin is scratched or rubbed. Neither papules nor vesicles are present in this form of the disease. The eruption is usually limited in extent, although it may occur acutely in the form of numerous small, rounded or oval patches scattered over the face, trunk, and extremities (Figs. 21 and 22).

In the erythematous eczema of childhood itching is not apt to be severe, although in adults this form occurring upon the



FIG. 23.—Eczema rubrum.

forehead, about the eyes, and elsewhere is apt to produce considerable thickening of the skin and a most intolerable pruritus.

In papular eczema, which was formerly described under the head of lichen simplex, we find irregular groups of bright-red, acuminate papules, many of which become excoriated and tipped by a minute blood crust, and some of which tend to run together, when sufficiently numerous, and form a thickened, scaly, or crusted patch. The accompanying plate shows a well-marked case of this form of the disease.

Vesicular eczema is a form rarely met with, since the natu-

ral duration of a vesicle rarely extends over a few days, and the itching which accompanies the eruption is certain to cause the lesions to be torn and the surface to become crusted by the drying of the serous exudation. And yet this aggregation of numerous minute, rounded, glistening vesicles upon a red and swollen surface of skin is perhaps the most characteristic phase of the disease. It was formerly believed that this was essentially a vesicular disease and that these lesions must be



FIG. 24.—Eczema impetiginosum.

present at the outset in every case of true eczema. The vesicular stage, when present, is attended with great heat and burning. The vesicles quickly rupture or coalesce, the epidermis cracks, and a surface exudation of a clear serum takes place. This is of a mucilaginous character, stiffens hair or linen with which it comes in contact, and dries like a varnish upon the inflamed surface, which quickly cracks, however, and thus allows the discharge to ooze out at many points. When the inflammation is very severe the greater portion of



the inflamed surface, instead of being crusted, appears swollen, moist, and of an angry red hue (*eczema rubrum*) (Fig. 23).

Pustular eczema results from an inflamed surface of skin becoming infected by pyogenic cocci. The characteristic oozing or "weeping" surface may gradually become purulent, and the exudation may dry into an extensive, thick, yellowish or honey-like crust (*eczema impetiginosum*) (Fig. 24); or a group of isolated follicular pustules or minute superficial abscesses may develop and become crusted, in which case the



FIG. 25.—*Eczema pustulosum*.

eruption bears a strong resemblance to contagious impetigo. This form of the disease is especially liable to occur in children of a strumous habit and among those who are not distinguished for their personal cleanliness (Fig. 25).

The squamous form of eczema is secondary to one of the preceding types. Indeed, a more or less scaly condition of the skin is seen in every patch of eczema before recovery takes place, and this may be regarded as the final stage in the course of the eruption. The amount and persistence of the scaling depend upon the extent of the infiltration of the skin. The

desquamation in eczema is always slight as compared with psoriasis, and the margin of the scaly patch usually shades off gradually into the surrounding healthy skin, instead of being circumscribed and abrupt as is always the case in the latter disease.

In the eczema of infancy, which is a very common and often a very obstinate affection, the face and scalp are usually affected, and frequently portions of the trunk and extremities suffer to a less degree. The characteristic tendency of the eruption to assume the moist form usually manifests itself, and the face appears either inflamed and angry or covered by dried exudation in the form of yellowish or blackish crusts. The little patient often suffers intensely from the oft-recurring pruritus, rolling his head in vain attempts to alleviate it by rubbing the inflamed skin against his clothing, or, if old enough to use his hands, tearing the face and causing it to bleed by sudden, fierce, and unexpected dabs with his sharp finger nails.

The cause of eczema might be discussed at great length and still little light thrown upon the subject, for it is certainly an obscure one. It may suffice to assert that improper food and imperfect digestion are the most frequent causes of the affection as it is met with in infancy.

Infantile eczema has usually nothing to do with the teeth. It often appears before teething begins, and a similar eruption may occur in old age when all the teeth are gone. Furthermore, it is always amenable to proper treatment, whether the child has one or twenty teeth. When the process of cutting teeth interferes with the general health of an infant, it is possible for the eruption to be indirectly aggravated by the condition of the gums, but in no case is "teething" ever to be regarded as the sole cause of eczema.

The "coming-out" of eczema is never a source of thankfulness, unless it happens to come out on some other baby than our own. We might be thankful if it were to come out in the baby's clothing, and it is about as likely to do this as it is to affect any of the internal organs. Eczema is essentially a disease of the skin, and cannot be transferred to the brain or lungs any more than the ruddy glow of a healthy cheek can be "driven in." It is true that sometimes an ointment is applied to an eczematous scalp or face, the eruption quickly disappears, and shortly after the baby may die of brain or lung disease. This may appear at first thought as though the eruption had been driven in, but the facts admit of a more reasonable



explanation. When from exposure to cold or some other cause an inflammation of the brain or lungs results, the blood flows in an unusual quantity to the affected organ, and any eruption upon the skin soon begins to fade and disappear. This, of course, takes place whether any ointment be applied or not. In such a case it might be said, with perhaps more truth, that the eruption has been drawn in, but it is evident that the disappearance of the skin disease is the result and not the cause of the internal inflammation.

The common belief that a skin disease is the outlet of some poisonous or vicious material in the blood is quite erroneous, and, in the light of modern pathology, a most absurd idea. An eruption can never be justly considered as either useful or beneficial to a patient. When a physician is unable to cure a case of infantile eczema, it may serve a purpose to claim that the eruption is salutary and that its sudden removal by local applications would be liable to produce disastrous results. The truth, however, is as follows: An eczema ought always to be cured as speedily as possible. This can always be done without the slightest danger of any harm to other organs of the body. Physicians of the largest experience in the treatment of skin diseases will all agree upon this point, and the sooner the old idea of the metastasis or translation of eczema is given up, the better it will be for the health of future babies and for the comfort of those who have the care of them.

In the treatment of eczema in infancy and childhood, it is well to remember that the disease is inflammatory in character, usually acute or subacute, and that our main object should therefore be to soothe the congested skin. If there were a law in this country prohibiting the use of any ointment save the officinal unguentum zinci oxidi, what a blessing it would be to children with acute eczema! As it is now, the physician is usually disposed to regard zinc ointment as little better than a domestic remedy, it being so well known. Familiarity with it has bred contempt. It will not do, he argues, for a man of his experience and reputation to prescribe so common and simple a remedy, and so he adds to it a little ichthyol and a little resorcin and a little carbolic acid, or possibly some one of the "new remedies" with which the general practitioner is usually familiar long before the specialist is willing to try them. The result is that the druggist is called upon to rub up a salve in which the bland, emollient character of the simple zinc oint-

ment is completely lost—a salve which is certain to irritate the delicate and inflamed skin and to aggravate the eruption.

More than a quarter-century ago Erasmus Wilson decided, after years of experience, that in acute infantile eczema the best method of treatment was by laxative doses of calomel and the external use of zinc ointment. While not unmindful of the progress which dermatology has made during this time, and in face of the flood of new remedies which is constantly pouring into our drug market, I defy any one in dermatological or pediatric practice to lay down a general method of treatment which, in the average case of infantile eczema, is practically superior to the one suggested by Wilson. A little starch or talcum added to zinc ointment will stiffen it and cause it to dry upon the skin, and thus obviate the necessity of smearing cloths and holding them in place by a bandage or mask. When a patch of eczema has ceased to exude and is in the final or squamous stage, a little oil of cade (one to five per cent) added to the zinc ointment or paste will greatly increase its efficacy in restoring the skin to a normal condition.

The tendency of soap and water to aggravate a moist eczema and to nullify the best of treatment is now generally understood by the profession. Indeed, the knowledge of this fact often leads the physician to forbid bathing in cases of dry eczema, when a daily bath would tend to improve the condition of the skin and benefit the patient. It is only in acutely inflamed and exuding eczema that water is necessarily injurious.

The regulation of the diet in case of eczematous children is of the utmost importance, as here may generally be discovered the cause of the eruption. With children old enough to go to the table, and especially with those who are apt to get whatever they cry for, a restriction of the diet to pure milk will often do much toward effecting a cure. It is very difficult to lay down dietetic rules of general application, for what will prove best suited to some children will fail to agree with many others; but it is certain that whenever a judicious local treatment appears to have little effect upon the eruption, the closest attention to the diet and condition of the stools becomes imperative. In many cases the local treatment which has produced no beneficial effect will work like magic as soon as the bowels are freely opened and the digestion improved.

The administration of arsenic, antimony, and other powerful

drugs may possibly do good in certain cases, but as I have seen arsenic, at least, do harm in a score of cases for every one in which I have seen it do good, I have no hesitancy in dispensing with their use in treating eczema in infants and young children. Arsenic should certainly never be given in the acute eczema of childhood nor during the exacerbations of the chronic form.





*PAPILLOMA LINEARE.*

From the collection of photographs of  
Dr. GEORGE HENRY FOX.



## PAPILLOMA LINEARE.

---

THE affection of the skin to which attention is now briefly called under the name of linear papilloma is somewhat rare and is described in the text books of dermatology under a variety of titles. It is frequently congenital, often develops in childhood, but may first appear in later life. It is characterized by warty excrescences, either colorless or pigmented, occurring in small groups running in one or more lines for some distance over the surface of the skin and frequently appearing upon only one side of the body. The neck, trunk, or one of the extremities may be the seat of the disease, and in rare cases the greater portion of the body may be affected. In congenital cases the warty outgrowth is apt to be fibrous and tough, while in cases developing later in life the lesions are softer, sometimes slightly scaly, and far more amenable to treatment. In some cases they have been known to disappear spontaneously or after some cutaneous exanthem, but a tendency to reappear upon the same site is usually manifested.

Of the names which have been applied to the affection the following may be mentioned: *ichthyosis hystrix seu localis*, *ichthyosis linearis neuropathica*, *nævus verrucosus*, *nævus unius lateris*, *nerve nævus*, and *papilloma neuroticum*.

The dark, fissured masses sometimes observed about the elbows and knees in severe cases of *ichthyosis* bear some resemblance to the warty lines seen in this affection, but they are simply accumulations of epidermis and not papillomatous in character. In linear papilloma, however extensive it may be, there is nothing suggestive of ordinary *ichthyosis*, and the statement made by some writers that a xerodermatous or parchment-like condition of the skin often coexists with papillomatous streaks is by no means in accord with my experience. The disease in question might be classed with ordinary *verruca* (warts) more justly than with *ichthyosis*.

The term *nævus verrucosus* would be applicable to this affection, but the fact that this name is commonly applied to the soft, flattened, pigmented, warty outgrowths so frequently seen upon the backs of old men and women renders its use objectionable in case of the disease under consideration. *Nævus unius lateris* is descriptive of most cases of this affection, but the fact that it may occur on both sides in certain cases may cause the name to sound absurd at times.



FIG. 26.—Linear papilloma.

The papillomatous streaks of this disease (like *zoster*) run transversely upon the trunk and longitudinally upon the extremities. They run in the direction of the natural cleavage lines of the skin, and, though they usually appear to follow the cutaneous distribution of certain nerves, they do not always do this with any great degree of accuracy. It would seem, therefore, that the name of linear papilloma as a descriptive title was preferable to the term nerve *nævus* or papilloma neuroticum.

In the accompanying plate, illustrating a case presented to the New York Dermatological Society by Dr. Cutler, the linear character of the affection is plainly shown in the streak coursing down the thigh and leg. In Fig. 26 the verrucous surface of the patch is quite apparent, while in Fig. 27 is seen the tendency of the lesions to form a broad, pigmented, warty patch in the axilla, from which a single line runs down the inner surface of the arm.



FIG. 27.—Linear papilloma.

The treatment of linear papilloma consists in the use of agents which tend to destroy or remove the excrescences. In cases of recent development, where the lesions are soft, the repeated application of a saturated solution of salicylic acid in collodion will act as effectively as it usually does in the case of corns and warts, and will soon leave the affected skin in a normal condition. In cases of longer standing it is advisable to use the curette, and where the eruption is extensive to remove

small portions of it at a time. But in certain cases, especially those which have existed since birth, the lesions are apt to be of such a dense fibrous character that the curette is of little use, and their removal is best effected by means of the curved scissors. It is doubtful whether any internal remedy is capable of affecting the growth in any notable degree.





Plate VIII.



Nævus pilosus.

From the collection of Photographs of Dr. George Henry Fox.

## NÆVI, PIGMENTED AND HAIRY.

---

THE term *nævus* signifies a spot or blemish upon the skin. It is usually a congenital affection, although certain forms of skin disease to which the term *nævus* is commonly applied may occur in youth or later in life. Many *nævi*, especially of the pigmented and hairy variety, exist at birth, but are almost imperceptible, and only become noticeable when they develop, as they are apt to do, after the age of puberty.

The term *nævus* includes affections of widely different pathological character in which hypertrophy is the only common element. They may be classed as fibrous, pigmented, hairy, and vascular *nævi*. The fibrous or hypertrophic *nævus* frequently occurring upon the face in the form of a small, rounded, and non-pigmented tumor or mole is peculiar to adult life, but the pigmented and hairy forms of *nævus* are common in childhood and often require treatment at an early age.

*Nævus pigmentosus* is the result of an excessive deposit of pigment granules in the mucous layer of the skin. It differs from a freckle or chloasmic discoloration (which are acquired affections and liable to disappear spontaneously) in appearing at or soon after birth and being permanent. The pigmentary *nævus* varies greatly in color and size. It may be yellowish or of a deep-brownish hue. It may be no larger than a pin's head, in which case it is apt to be multiple, or it may cover a portion of the body larger than the hand. The face, neck, and back of the hand are favorite sites, but it may appear on the trunk and extremities. It is sometimes flat and smooth, like a freckle, but may become elevated and warty in certain cases, when the term *nævus verrucosus* is applicable.

In many cases of extensive pigmentary *nævus*, whether smooth or elevated, we find a growth of fine or coarse hair upon the surface. When this is more marked than the pigmentation the affection is called *nævus pilosus*. The hairy *nævus* is often

a small oval patch (as in the plate), and from its size and shape its origin is frequently ascribed to some maternal impression—*e.g.*, the mother having been frightened by a mouse. Often the hairy scalp appears to extend over the temple or upon one side of the forehead, as in Fig. 28, and not infrequently it is seen



FIG. 28.

upon the cheek beneath the eye or involving both lids and vicinity, as in Fig. 29.

In rare cases a large extent of cutaneous surface is affected, and the whole trunk may appear as though covered with fur instead of normal skin (Fig. 30). The lumbar or pelvic region is also apt to be the seat of the hairy nævus, which in some cases

has suggested a resemblance to "bathing-tights." The congenital growth rarely increases in extent, except as the body grows larger, but often the pigmentation becomes more pronounced and the hair begins to grow coarser in adult life. The development of small hairy moles upon the face of elderly



FIG. 29.

women, especially those suffering from hypertrichosis, is very frequently noted.

In the treatment of the pigmentary mole, acids or the electrolytic needle may be employed. For small, round, dark spots on the skin a minute drop of nitric acid, applied with a wooden toothpick, will often suffice to remove the blemish. In case of larger pigmented patches, either smooth or covered with fine hair, the surface of the skin may be dotted with the acid; but



extreme caution must be used to prevent ulceration, lest scars be left which would be far more disfiguring than the nævus itself. With children old enough to bear a little pain without crying, the pigmented spots may be blistered by touching them with the point of the electrolytic needle or by passing it superficially through the epidermis.

In nævi of small or moderate size covered with coarse hair



FIG. 30.

the electrolytic needle may be used, as in the treatment of hypertrichosis, and a fine result attained through patience and perseverance. In the *Journal of Cutaneous Diseases* for May, 1893, the writer has reported a case of extensive hairy and verrucous nævus involving the right cheek and lower eyelid, which was completely removed by means of electrolysis. The treatment consisted in carefully passing a fine, flexible steel



needle, connected with the negative pole of a galvanic battery, through the most superficial portion of the growth, the circuit being completed by the patient grasping a moist sponge attached to the positive electrode. This was repeated until the electrolytic destruction of tissue reduced the growth to the level of the surrounding skin, removed the pigmentation, and to a certain extent destroyed the hypertrophied hair follicles. The slight growth of hair which persisted after the affected skin had become smooth and comparatively normal in color was destroyed by the introduction of the electrolytic needle into each separate follicle, according to the method employed in the treatment of superfluous hair. The young man now presents not the slightest deformity of the eyelid, and scarcely a trace, upon close inspection, of the dark, warty, and hairy growth which formerly attracted attention wherever he went and was the source of great mortification.





PLATE IX.



Vascular Nævus.

From the collection of Photographs of Dr. George Henry Fox

## VASCULAR NÆVUS.

---

THE vascular form of nævus results from an excessive development of vascular tissue in or beneath the skin, and presents a variety of clinical appearances. These may be conveniently classified as follows : 1, nævus araneus ; 2, nævus flammeus, or port-wine mark ; 3, nævus tuberosus ; and 4, nævus cavernosus. Of the many descriptive names which have been more or less in use, these appear to best express the characteristic features of the four clinical varieties of vascular nævus.

Nævus araneus consists of a small red spot, smooth or slightly elevated, from which several dilated capillaries radiate like the legs of a spider. It has sometimes been called "spider cancer," a name which is as alarming as it is inappropriate. It is not always congenital, but frequently appears in youth or adult life as the result of a prick, bruise, or other injury to the skin. On this account some writers class it under the head of telangiectasis or acquired vascular dilatation, assuming that the term nævus signifies a birthmark and not merely a spot or blemish. It is most frequently noted upon the upper portion of the face, and upon a fair white skin may be quite conspicuous and annoying.

Nævus flammeus, birthmark or port-wine mark, as it is commonly called, usually appears as a smooth, red discoloration upon one side of the face. It varies greatly in size, often extending down upon the neck and involving a considerable portion of the trunk. It varies also in color from a light pink to a deep purplish-red or even slate-colored hue. In its severe form the affected skin is elevated, the lip greatly swollen, and one side of the face may appear notably larger than the other. The dark surface of the skin may be dotted here and there in some cases by small vascular excrescences. At birth small, faint red marks may be noted, especially upon the lower portion of the occiput, which tend to disappear in later years, but the port-wine mark usually persists throughout life, either remaining unchanged or becoming more conspicuous. It has been claimed



that some cases have increased in area and that others have gradually disappeared, leaving atrophic scars, but such cases are notable exceptions to the rule.

Nævus tuberosus is a turgescient tumor of varying size with a rounded or flattened surface. It results from a congenital dilatation or new formation of blood vessels, and in color varies from a bright cherry-red to a purplish-red or leaden hue, according to the predominance of arterial or venous blood in the



FIG 31.—Nævus of nose.

tumor. It is frequently seen upon the scalp and face, although it may be found upon the trunk and extremities. It usually increases somewhat in size during the early months of infancy, and in some cases develops with such alarming rapidity that vigorous treatment is called for without delay. The tumor is usually elastic and compressible, and often varies in size and color at different times of day, and is especially prominent after a fit of crying or severe coughing. A distinct pulsation may

be sometimes felt, corresponding in rhythm with the action of the heart. Ulceration of the central portion of the surface occasionally takes place, and from this cause or from some accidental injury severe hemorrhage may result. Gangrene may occur and produce a spontaneous cure.

Nævus cavernosus is a deep-seated form of angioma over which the raised skin may appear of normal hue or present a dull-bluish or venous appearance. The tumor is formed by



FIG. 32.—Birthmark of unusual extent.

masses of dilated veins and arteries surrounded by firm connective tissue, which extends into the interior and forms vascular cavities communicating freely with the enlarged vessels. In some cases a number of these tumors, of varying size but of the same character, will extend along the surface of the skin like deep varices. They are usually soft, especially when of large size, and have a peculiar lobulated feeling when pressed beneath the fingers.

In the treatment of *nævus araneus* and other telangiectases the electrolytic needle is most serviceable. Any form of galvanic battery may be used, and the point of an ordinary cambric needle or flexible steel jeweller's broach, attached to the negative cord, should be pressed into the centre of the red spot. A moist sponge attached to the positive cord should now be grasped in the patient's hand or slowly applied to the skin at any point, when the electrolytic current will begin to act and quickly produce a whitening of the skin around the needle



FIG. 33.—*Nævus* of scalp.

point and cause sufficient inflammation to seal up the deep-seated supply vessel. The pain caused by this operation is slight and no more than the promise of some candy or a coveted toy will induce the average child to endure. The result is always excellent.

The electrolytic needle may also be used with success in many cases of port-wine mark, but the treatment is tedious and only capable of producing good results when persistently and skilfully employed. The best that can be said of it is that in



case of extensive and dark-hued patches it is superior to any other plan of treatment, even if it is not productive of the speedy and brilliant result which might be desired. The object of this method of treatment is to cover the dark-red skin with minute punctate cicatrices, which will at least lessen the conspicuous character of the mark, if it does not remove it entirely. To remove a red birthmark and leave a perfectly normal skin is an impossibility. In cases where the color is not



FIG. 34. —Nævus tuberosus.

very deep a good result may be obtained by dotting the surface with minute drops of nitric acid, great care being taken that these do not spread or run together and in this way produce ulceration and subsequent pitted or raised cicatrices.

In the tuberoso variety of nævus, where red tumors suggestive of a strawberry, cherry, or flattened tomato are to be removed, a resulting cicatrix is unavoidable, and either acids, electrolysis, or the platinum cautery may be advantageously used. The size and disfiguring character of the scar which is

necessarily produced will depend in great measure, however, upon the skill and caution displayed in the treatment, whatever means may be employed in destroying the vascular growth. The attempt to remove this form of *nævus* by compression has always failed, in my experience; and as for ethylate of sodium,



FIG. 35.—*Nævus* of vulva.

although it has proved successful, it has been quite as painful and less effective than nitric acid.

In the treatment of the cavernous *nævus* the knife, ligature, galvano-cautery, and injections of carbolic acid and of iodine have been recommended, and the most suitable method of treatment must depend upon the nature of the case.





PLATE X.



Lapus Vulgaris.

From the collection of Photographs of Dr George Henry Fox.

## LUPUS AND OTHER TUBERCULIDES.

---

MANY years ago the fact was noted by dermatological clinicians that lupus vulgaris was frequently associated with pul-



FIG. 36.—A group of tubercles.

monary phthisis, and recent microscopical research has demonstrated that in both affections the same bacillus may be found. Lupus vulgaris is therefore one of the clinical forms of cutane-

ous tuberculosis and may be properly described as a tuberculide. There are other affections of the skin which are commonly associated with the varied symptoms of scrofula and in which the bacillus tuberculosis may be found, but they are clinically distinct, though pathologically related to lupus. The term tubercular, as applied to lupus, was originally used only in a morphological sense, indicating the nodular character of the eruption. It is a singular fact that, while the old term is still



FIG. 37.—Tubercles forming a ring.

retained in use, science has invested it with a deeper and more definite significance since the discovery in the nodule of the tubercle bacillus.

Lupus vulgaris is an affection which is often seen in childhood, and which, indeed, begins in most cases before the age of puberty. It usually appears in the form of one or more dull red papules upon the cheek or elsewhere. These slowly increase in number and tend to coalesce (Fig. 36). Frequently the older or central lesions disappear by interstitial absorption and an

irregular ring is formed (Fig. 37), or an infiltrated patch with one or two outlying nodules (see plate). Upon the neck the disease often assumes a serpiginous form, spreading at the margin and enclosing a cicatricial area, dotted, perhaps, by a few recently developed nodules (Fig. 38). This form of the disease may have a somewhat acute character, and, though spreading slowly, may increase in extent much more rapidly than does the nodular form seen upon the cheeks.

Upon the ala nasi, which is a frequent site of lupus, a scaly



FIG. 38.—Serpiginous lupus with central cicatrix.

or crusted patch is often seen, with more or less ulceration, and in time a marked deformity as the result of cicatricial contraction (Fig. 39).

The nodules of lupus, when well developed, have a characteristic translucent, jelly-like appearance, and, though feeling firm and resilient to the touch, are much softer than the normal cutaneous tissue. In an advanced stage the nodules and patches of lupus become somewhat scaly and not infrequently soften and ulcerate. In chronic cases seen in adult life the



disease is often found to have spread over the greater portion of the face, producing a marked ectropion and a partial disappearance of the nose and ears. Such an extensive development of the disease is rarely if ever observed in childhood. Although the face is the most common site of lupus, the trunk and extremities may also be affected, either independently or with the face.



FIG. 39.—A favorite site of lupus.

Another tuberculide, or affection in which the bacillus tuberculosis is invariably present, is often met with in childhood as well as in adult life, and is known as tuberculosis verrucosa or lupus verrucosus. This consists of dry, warty, or papillomatous patches (Fig. 40), which may develop upon the back of the hand, especially over the knuckles, the wrist, the popliteal spaces, and other portions of the body, and is commonly regarded as the result of local infection. The development of the disease is

slow. Ulceration rarely occurs, but the centre of the patch may undergo a spontaneous cure, leaving cicatricial tissue.

Still another form of cutaneous tuberculosis is that commonly described under the name of scrofuloderma. This appears in the form of suppurating or crusted ulcers of the skin in children who usually present other evidences of the scrofulous taint. It is most frequently observed upon the neck over




FIG. 40.—Lupus verrucosus (tuberculosis verrucosa cutis).

lymphatic glands which have undergone caseous degeneration and softening, although it may occur in multiple discoid lesions over the buttocks, thighs, and other portions of the body. When lupus or scrofuloderma attacks the hand or fingers of a child the bony tissue is frequently involved and a strumous dactylitis may develop, or caries with resulting atrophy and a considerable subsequent deformity (Fig. 41).

Lupus erythematosus, though allied in name to lupus vulgaris and often bearing a strong resemblance to it, cannot be considered as a tuberculide. The bacillus tuberculosis has not been found in this affection, and all attempts at inoculation of animals have given negative results. It may be remarked in passing that this disease, like lupus vulgaris, is most commonly seen upon the face (Fig. 42), but, unlike the common form of lupus, it is rarely seen in childhood.



FIG. 41.—Ulcerating lupus with atrophy of bone.

In the treatment of lupus and the other tuberculides much may be done to improve the general health of the patient and thus to modify to a limited extent the spread of the disease; but to effect a cure a resort to surgical measures is necessary. The palliative treatment by means of ointments and plasters, so frequently employed, need only be mentioned for the sake of condemnation. The morbid tissue must be destroyed, and it matters little how this is done provided it is thoroughly done.



The knife, the cautery, and caustic pastes I have used and discarded, believing that by the skilful use of the curette and burr the disease can be removed with the least amount of pain and discomfort and the least resulting disfigurement. For many years I have used the dental burr of varying size, dipped in carbolic acid, for the destruction of lupus nodules, and ever with increasing satisfaction. This instrument readily penetrates the



FIG. 42.—Lupus erythematosus.

gelatinous lesions, and when the handle is rolled between the fingers, and the burr pressed in various directions, it bores out the softened lupus tissue as it does the carious substance in a dental cavity and leaves the normal skin uninjured. In diffused patches of lupus, in scrofulous ulcerations, and in verrucous tuberculosis (after the warty surface has been removed by a salicylic-acid plaster) nothing can be more serviceable than the dermal curette.







Plate XI



PHOTOGRAPH BY S. CULLEN OF N.Y.

LICHEN RUBER

(*Pityriasis rubra pilaris*)

*From the collection of photographs of Dr. George Henry Fox*

## LICHEN RUBER AND LICHEN PLANUS.

---

LICHEN RUBER, or pityriasis rubra pilaris as it is called by many writers, is a rare disease, but one which is liable to affect children as well as adults. It is a chronic and obstinate disease, in which the lesions are small, reddish, acuminate papules,



FIG. 43.—Papular form of lichen ruber.

usually seated at the orifice of hair follicles. These do not increase in size, but by multiplying and coalescing produce patches covered by fine, whitish scales and characterized by an exaggeration of the natural furrows of the skin.

The eruption often undergoes a spontaneous improvement,




FIG. 44.—Papules on a favorite site.



FIG. 45.—Plantar lichen ruber resembling eczema.

but only to relapse, and, as a consequence of this, its clinical



appearance varies greatly at different times and often upon




FIG. 46.—Lichen ruber resembling ichthyosis.



FIG. 47.—Lichen ruber resembling psoriasis.

various portions of the body. Three stages or clinical forms



of the eruption may be conveniently described as the papular, squamous, and rugous forms of the disease.

In the papular form of lichen ruber (Figs. 43 and 44) the lesions are small, usually acuminate, and often tipped with a minute white scale. When of recent development they produce an appearance quite similar to *cutis anserina* or follicular keratosis. Upon the dorsum of the fingers there is often noted a group of follicles plugged with blackened epidermic scales, while upon the palms and soles a scaly condition is usually pres-




FIG. 48.—Rugous form of lichen ruber.

ent, closely resembling a chronic squamous eczema (Fig. 45). The papules often increase rapidly in number upon certain portions of the body, become flattened and scaly as they coalesce, and now present an entirely different aspect.

In the squamous form of lichen ruber the eruption appears as white, scaly patches of varying size and form, and frequently bears a strong resemblance to ichthyosis (Fig. 46) or psoriasis (Fig. 47). The eruption in this stage shows a marked tendency to occur in elongated, spindle-shaped, or band-like patches.

At the margin of these squamous patches numerous isolated, white-tipped papular lesions may frequently be seen. The scalp is generally affected when the disease has existed for some time, and the face often presents an appearance suggestive of ichthyosis.

In the rugous form or stage of the disease the affected portions of skin present a dull-red, leathery appearance, with slight scaling and deep parallel furrows (see plate and Fig. 48). Upon the hands the skin sometimes becomes shrivelled and drawn to an extent which seriously interferes with the motion of the fingers (Fig. 49), while the nails show marked evidence of malnutrition. In some cases nodular ridges are found at the bend



FIG. 49.—Chronic form of disease.

of the elbow, upon the pubes, and elsewhere, and present a moniliform or bead-like appearance (Fig. 50).

Itching is commonly present, and often very severe and annoying in advanced cases, and an exacerbation of the eruption is frequently preceded by an intense burning sensation.

The prognosis in lichen ruber is an unfavorable one. Although the disease may yield to judicious treatment, and even improve spontaneously, a long series of relapses may be looked for, and a fatal termination be predicted in many cases. The reported cure of cases by French and German writers may be readily explained by their belief in the identity of lichen ruber and lichen planus.

From what has been said it is evident that the treatment of this disease must of necessity be palliative rather than curative. The improvement of the patient's general health by a hygienic and tonic regimen will usually accomplish more than the administration of arsenic or any other drug. Meanwhile the comfort of the patient can be greatly increased by baths and inunctions tending to soften the dry, harsh skin and to lessen the pruritus.

Lichen planus is a disease which is entirely distinct in nature



FIG. 50.—Moniliform lichen ruber.

from lichen ruber, although the two affections have been considered as clinical forms of the same disease by many European writers. This erroneous view has led to considerable confusion of dermatological literature. The typical lesions of lichen planus are small, flattened, angular papules with a shining surface and a minute central depression. Those of lichen ruber, on the other hand, are usually acuminate, although in rare cases they may appear flattened and smooth. The eruption in lichen planus commonly presents a purplish or lilac hue, which is very

characteristic and often serves as an excellent basis of diagnosis. The lesions are at first discrete, but show a marked tendency to coalesce and form irregular or reticulate patches of varying




FIG. 51.—Lichen planus.

size. The most common site of the eruption is upon the anterior aspect of the forearm and next upon the lower extremities. It is often seen around the waist and sometimes upon the



FIG. 52.—Lichen planus.

genitals. In exceptional cases the greater portion of the trunk may be affected, and such cases are very apt to be confounded with lichen ruber. Upon the legs the patches are frequently



roughened and pigmented, and present an appearance quite unlike the eruption seen upon the trunk and forearms.

Lichen planus runs a variable course, some cases disappearing spontaneously and often unexpectedly after an existence of a month or two, while others will sometimes persist for many months in spite of the most approved method of treatment. The itching is often very annoying to the patient, but the general health is usually unimpaired, and, unlike lichen ruber, the disease never terminates fatally.




FIG. 53.—Lichen planus.

In the treatment of lichen planus arsenic is often of service, but in this, as in many other affections, it may do harm as well as good ; and when there is much irritability of the skin alkaline remedies will be found to be of much greater service. Of the various local remedies which have been highly recommended no one seems to have any special value in all cases. A mild carbolie or salicylic acid lotion may be advantageously used in acute cases, and the same increased in strength when the patches have assumed a chronic character.





Plate XII



PHOTOGRAVURE & COLOR CO. N. Y.

SYPHILIS HEREDITARIA

*From the collection of photographs of*

*Dr. George Henry Fox.*

## VARIOUS SKIN DISEASES OF CHILDREN.

---

*Syphilis hereditaria*.—When syphilis is transmitted from one generation to another it may result in the death of the foetus in utero, or in the development of an infantile eruption which either is present at birth or appears shortly after. In most cases it manifests itself within three months, and always during the first year. The eruption may be erythematous and appear as a general roseola, similar to the earliest eruption of acquired syphilis, or in the form of dull-red patches on the trunk and upon the palms and soles, which present a dry, scaly appearance quite characteristic of the inherited disease. About the mouth the skin is often dry and fissured, and on the buttocks the patches are apt to become raw and even ulcerated at various points. In some cases the eruption assumes a papular character (see plate), and numerous yellowish-red and slightly scaly discs are seen upon the buttocks, thighs, and face, and occasionally over the greater portion of the body. In rare instances the papules are small, flattened, and shining, and present a strong resemblance to lichen planus. Pustular lesions of an ecthymatous character are sometimes present, especially in infants who are ill-nourished and poorly cared for. Bullous lesions are not infrequently seen upon the palms and soles and other portions of the body, and in fact are much more common than in the acquired disease of adult life.

In connection with these early cutaneous manifestations a notable condition of malnutrition is usually observed, and certain characteristic symptoms resulting from affection of the mucous membranes and the osseous structures. The nasal passages are often inflamed and obstructed by an accumulation of mucus. The larynx may be affected, giving rise to a peculiar hoarse cry.

The prevention of hereditary syphilis is more important than

its treatment, and the physician is often in a position to advise against the premature marriage of syphilitic patients, and thus prevent an amount of misery and suffering which he could never cure.

In the treatment of infantile syphilis a half-grain or more of mercury with chalk may be administered three times a day, and a speedy improvement expected if the child is properly cared for. When irritability of the stomach and intestines is present it is advisable to adopt the plan of mercurial inunctions. A mass of mercurial ointment, the size of a small marble, should be spread upon a piece of flannel and bound around the trunk or thighs, its position changed from day to day to prevent excessive irritation of the skin.

In treating mucous tubercles and superficial ulceration upon the buttocks and about the genitals, calomel powder mixed with three to five parts of starch may be sprinkled over the raw surface, care being taken to keep the affected parts dry and clean. The importance of fresh air and nutritious food must never be forgotten, and when the symptoms of the disease have abated the syrup of iodide of iron, and perhaps cod-liver oil, may usually be given with great benefit.

*Erythema nodosum* is one of the forms of erythema multiforme, though commonly described in dermatological text books as a distinct disease. It occurs most frequently in childhood, and the female sex seems especially prone to suffer from it.

The lesions in this affection are red, painful swellings, looking very much like recent contusions, and are most frequently seen over the tibiæ. They are usually associated with slight fever, considerable malaise, and in some cases with articular pains. Both legs are commonly affected, and rarely are the lesions found elsewhere. In some cases, however, the lesions are numerous and found upon the thighs as well as the legs (Fig. 54), and in rare instances the forearms may be affected. The lesions are at first of a bright-red hue, but soon become dull red or purplish, and usually disappear in a week or two. New lesions sometimes continue to appear and prolong the duration of the disease. The individual nodules usually last from eight to fourteen days. New nodules appear, as a rule, during the first two weeks, and the disease may therefore last from three to six weeks. In exceptional cases it may continue for months.

The cause of erythema nodosum is obscure. Most patients appear to be in poor condition, and a rheumatic tendency is



sometimes noted. An examination of the blood will sometimes reveal the presence of malarial organisms.

The eruption always tends to a spontaneous recovery and usually shows no disposition to recur. While no medicinal treatment is absolutely necessary, a saline laxative at the outset will often do some good, and may be followed by the administration of salicylate of soda.

Locally a lead-and-opium wash, or a lotion of zinc oxide in lime water, will tend to allay the burning pain or tenderness of



FIG. 54.—Erythema nodosum.

the lesions, which is often a distressing symptom, and rest in bed for a few days is desirable during the height of the attack.

*Purpura*, or idiopathic hemorrhage into the cutaneous tissues, results from many causes and occurs in connection with various diseases. It is usually characterized by small petechial or larger guttate spots, which are at first of a bright claret hue, but which rapidly assume a dull-purplish color. The eruption is readily distinguishable from an erythema or any inflammatory exanthem, since the redness does not disappear under firm pressure of the finger.

In *purpura simplex* the lesions are commonly symmetrical and seen upon the legs, but in children more frequently than



in adults the eruption may appear upon the thighs (Fig. 55) and other portions of the body. In a child suffering from whooping cough I have seen the chest dotted with petechial spots. When the hemorrhage takes place in or around a hair follicle the lesions are apt to be elevated, and to this condition the term *purpura papulosa* has been applied.

In mild cases of *purpura simplex* the lesions usually appear suddenly without other symptoms, and gradually fade away



FIG. 55.—Purpura.

in one or two weeks, unless a fresh outbreak occurs. In this event the bright-red color of the recent spots contrasts strongly with the dull livid hue of the older lesions.

In severe cases of *purpura* the eruption may be very abundant, and with the cutaneous lesions hemorrhage from various mucous surfaces may occur. The gums are usually swollen and bleed easily. Epistaxis is frequent and blood is often noted in the stools and urine. This form of the disease is called *purpura hemorrhagica*. It is characterized by marked

constitutional symptoms, and the weakness resulting from a continued loss of blood sometimes leads to a fatal termination.

Another form of the disease is associated with severe arthritic pains and the appearance of elevated purpuric lesions, especially in the vicinity of the affected joints. This is termed purpura or peliosis rheumatica. This affection usually runs a brief course, although a repeated onset of fever and rheumatic pains, followed by cutaneous hemorrhages, may protract it for months. It commonly attacks those who have a rheumatic tendency but are otherwise apparently well.

There is little difficulty in the diagnosis of purpura, as the color of the lesions is characteristic and their hemorrhagic nature can be readily demonstrated by the pressure of the finger. It is well to remember, however, that cutaneous hemorrhage is not infrequently an accompaniment of the eruption in severe cases of the common exanthemata.

In the treatment of mild cases of purpura, rest in bed, with a nutritious diet and administration of the *tinctura ferri chloridi*, is generally followed by a prompt recovery. In severe cases, especially of the hemorrhagic type, iron, turpentine, and ergot are commonly employed with good effect. Ergot may be given by the mouth or by hypodermatic injection. For the latter purpose a grain of ergotin may be dissolved in warm water and glycerin, and injected with such frequency as the exigencies of the case may demand.

*Molluscum* is a term which has been applied to two distinct affections having no pathological relation to one another. One is the *molluscum fibrosum* of older writers; which is now commonly designated as fibroma. The other is the *molluscum contagiosum*, which was first described by the English dermatologist Bateman early in this century. The latter is a somewhat rare affection of the skin, but of much more frequent occurrence in childhood than in adult life. It consists of one or more small, flattened whitish tumors, varying in size from a pin's head at the outset to a split pea when fully developed. These little growths are sometimes pedunculated and usually present a central depression, from which a whitish substance can be readily pressed out. This looks like ordinary sebaceous matter, but under the microscope is found to contain certain oval corpuscles which are a characteristic of the disease.

The tumors are commonly seen upon the face and neck, but may occur on various portions of the body. They are usually multiple, sometimes numerous, and, though unsightly, are

never the source of any particular discomfort. They develop slowly and last indefinitely. Each begins as a minute elevation of a whitish color, and gradually the flattened summit and umbilication develops. When of full size the walls are sometimes traversed by fine blood vessels which give the tumor a pinkish appearance.

The disease is often called *molluscum contagiosum*, and, though its contagious character is not always apparent, it has



FIG. 56.—Molluscum.

been known to affect several in a family or in a ward of a children's hospital, and attempts at artificial inoculation have sometimes been successful. It is doubtless of microbic origin, but the conditions favoring its development are as yet obscure. It is especially liable to affect children of a poorer class, and hence is much more common in dispensary than in private practice. Damp and crowded dwellings seem to favor its development, and in dispensary practice I have known a number of



cases to originate in the same locality. Children suffering from molluscum are especially prone to have warts upon the hands.

The diagnosis of molluscum is usually made with ease by any one at all familiar with the appearance of the tumors. Large milia upon the face might be mistaken for them; but these tumors are rounded, of much firmer consistence, not likely to occur in groups, and always lack the characteristic flattened surface and central umbilication.

The treatment of molluscum is simple and effective. Many of the tumors, after a duration of a few weeks or months, undergo a process of destructive inflammation, and a spontaneous cure results. This is especially apt to be the case when two are close together and coalesce as they increase in size. If a tumor is vigorously compressed and the contents evacuated it will usually disappear. Incision and cauterization, or abscission by means of a sharp knife, have been recommended, but the simplest method of removing the tumors is to scrape them quickly with a curette and to lightly touch the bleeding surface with nitrate of silver. The main object in treating these tumors is to destroy them completely and to excite as little inflammation as possible in so doing. When even a large molluscum is removed by means of the curette no scar or trace of the growth is usually left, since the tumor is an outgrowth of the epidermis and does not involve the true skin. But when an inflamed molluscum is scratched or irritated to the extent of inducing a purulent secretion, a slight pit may be expected to result.

*Keratosis follicularis* is an affection not infrequently seen in childhood, especially among children who are poorly cared for. It is the result of an abnormally dry condition of the skin and the accumulation of epidermic scales in the hair follicles. This produces an eruption of small, conical papules, which are commonly found upon the outer aspect of the extremities. These are usually of the color of the skin. Sometimes they are discolored by dirt, and in some cases they become inflamed and present a dull-red hue. The fine hairs growing upon the affected skin are often broken off or coiled up in the follicles. When the disease is severe and of several years' duration the skin has an ichthyotic appearance and to the touch feels like a nutmeg grater. The disease is described in some text books under the name of lichen pilaris or keratosis pilaris, but the follicle is primarily involved and not the hair. Recently the term *keratosis follicularis* has been applied by some writers to

an entirely different and extremely rare disease described by Darier.

The treatment of the disease is mainly palliative, as in many cases time alone will effect a complete cure. The daily inunction of the skin with some fat or oil, and the frequent resort to hot baths and vigorous soap frictions, will be productive of much benefit and often restore the skin to its normal condition.



FIG. 57.—Keratosis follicularis.

But this treatment must be continued for some time to prevent a return of the disease. A Turkish bath once or twice a week will prove of service.

*Keloid* is a dense fibrous tumor of the skin, which in certain individuals is liable to develop upon the site of a cut, burn, or other injury. A distinction has often been made between true and false keloid, many writers claiming that the former develops spontaneously while the latter is always an outgrowth upon



a scar. This distinction, however, is of little value, since the growth is the same in either case, and it is impossible to assert, in any case of spontaneous keloid, that it has not been preceded by some prick or slight injury to the skin. A distinction, however, may be justly drawn between keloid and a hypertrophic cicatrix. Many scars, especially those following burns, are prone to pucker and bulge until a tumor is formed which looks very much like keloid, being rounded, smooth, reddened, and often very firm and even painful. But while keloid is characterized by a marked tendency to enlarge its area by the forma-



FIG. 58.—Keloid.

tion of claw-like processes suggestive of a crab, and to persist indefinitely, the hypertrophic cicatrix never extends beyond the limits of the scar tissue and tends to a gradual disappearance rather than to an increase in size.

Keloid is usually single, but many tumors may be present, and in some cases these attain considerable size. The growth is commonly painful when squeezed or pressed, and sometimes a continuous burning or pricking sensation is experienced. A spontaneous disappearance of the tumors has been reported, but this is exceptional. Keloid occurs at all ages, affects various

portions of the body in both sexes, and is especially common in the negro race. The tumors increase in size for a certain time and then often remain stationary for years. A characteristic peculiarity is their tendency to return quickly whenever excised. A spontaneous disappearance of keloid is more likely to occur in childhood than in adult life, and tumors developing upon large scars are more likely to disappear than those which seemingly spring from the normal skin. A favorite site of keloid is the sternal region, where it often appears in adults as an elongated flattened tumor crossing the median line in a transverse direction. Its frequent occurrence in this locality has been attributed to the common custom of applying blisters to this part for the relief of acute pulmonary affections. It also is seen frequently upon the scalp, face, neck, and trunk.

The treatment of keloid is usually discouraging. Excision, as has been remarked, is usually followed by a return of the growth. Good results have been obtained by deep linear scarification and the application of glacial acetic acid to the cuts, and also by the use of the electrolytic needle. Subcutaneous injections of thiosinamine have been recommended, and in hypertrophic cicatrix I have known this to be followed by benefit. The results from the subcutaneous use of this drug in lupus, psoriasis, and certain other skin diseases have not proved to be as beneficial as its enthusiastic advocates of a few years since were disposed to claim. But in the relief of corneal opacities and cicatricial contractions it has apparently proved of value. Whether its use in true keloid will effect even a notable improvement remains to be demonstrated; but in cases of scar keloid which are tending to recovery it might, at least, be tried in connection with the fatty inunctions and frequent massage which seem to do some good and certainly do no harm.

*Scabies* is a disease which in this country has notably increased in frequency during recent years. In most cases it is readily recognized by the characteristic location of the excoriated papules which mainly constitute the eruption. The disease, as is well known, is due to the burrowing of the acarus or itch-mite in the soft skin between the fingers and elsewhere, and is usually communicated by sleeping with, or in a bed which has been occupied by, a person affected with the disease. The mature, impregnated female alone makes the burrow in which the eggs are deposited, and dies after performing this function. In recent cases the burrows or cuniculi in which the female acarus has deposited her eggs may be found upon the web of the

fingers, the anterior portion of the wrist, the axillæ, genitals, and ankles, and appear as small, curving dotted lines, from the extremity of which the insect may sometimes be dug out with the point of a needle. These burrows are often concealed by an eruption of vesicles, pustules, and crusted sores, and not infrequently the lesions of contagious impetigo are present. Upon other portions of the body the eruption usually consists of numerous excoriated papules covered by minute blood crusts. The anterior aspect of the forearms and the abdomen are especially apt to be the seat of the eruption, but when the disease has existed for several months the eruption may cover the



FIG. 59.—Scabies.

whole body, with the exception of the face, which always remains unaffected, except in the case of infants and very young children. The location of the eruption in this disease depends to a great extent upon the character of the clothing which the patient wears, and differs in the case of men, women, and children. The excoriations about the breasts of women and upon the genitals of men and boys are due to the ease with which these parts can be scratched at night, while in young children who wear long, close-fitting night-dresses the wrists and ankles are most likely to be the seat of the eruption.

The treatment of scabies is simple and effective. For gene-



rations sulphur ointment has been a standard remedy, and nothing else is needed, although many other remedies and combinations have been recommended. A warm bath should first be taken to soften the skin, and soap used vigorously to remove the dead epidermis covering the cuniculi. The sulphur ointment should now be rubbed gently into the affected skin, especially in those regions where the acarus is apt to burrow,



FIG. 60.—Pustules in scabies.

and this inunction repeated, *without bathing*, for about five successive nights. Under this treatment the itching is abated and the disease usually cured. In severe cases it may be necessary at the end of five days to repeat the bath and the subsequent inunctions. In infants and young children, or in patients with a very delicate skin, it is advisable to dilute the sulphur ointment with one or two parts of vaseline or cold cream.

## FORMULARY.

---

NOTE.—In the following formulary the decimal system has been employed, and the percentage of each ingredient can therefore be readily seen at a glance. The unit may be regarded as a grain, a gramme, or any other weight, without changing the relative amount of the ingredients. An exception to this rule will be found in the baths recommended for syphilis, in which definite amounts are given in both the metric and the apothecaries' systems. The formulæ have been carefully selected from the writings of various well-known authorities, and some trifling changes have been made in a few cases in order to reproduce the prescriptions in a decimal form or to render them more suitable for use in childhood. The signs R, M., and S. have been omitted to avoid repetition. As far as possible an attempt has been made to give due credit to the author of each formula.

### ACNE.

1.

Glycerini.....	3
Alcoholis.....	20
Sulphuris præcipitati.....	30
Aquæ destillatæ.....	q. s. ad 100.

To be used as a lotion.

(*Liveing.*)

2.

Ichthyolis.....	10
Alcoholis.....	50
Ætheris.....	q. s. ad 100.

3.

Spiritus lavandulæ.....	3
Saponis mollis.....	6
Tincturæ hamamelidis.....	10
Alcoholis.....	q. s. ad 100.

Use to stimulate indurated spots.



## 4.

Acidi borici.....	3
Aquæ hamamelidis .....	40
Aquæ rosæ fortioris.....	q. s. ad 100.

To be used in acne punctata.

(Shoemaker.)

## 5.

Iodi ...	25
Potassii iodidi.....	25
Glycerini.....	q. s. ad 100.

Paint the surface with a brush twice daily for from three to six days. When the crust is cast off apply a simple ointment or powder. This is to be used in severe indurated cases only.

## 6.

Tincturæ benzoini.....	2
Glycerini.....	3
Zinci oxidi.....	10
Unguenti aquæ rosæ.....	q. s. ad 100.

## 7.

Thymolis.....	1
Oleati zinci.....	6
Sulphuris sublimati....	12
Adipis lanæ hydrosi.....	q. s. ad 100.

## 8.

Europheii .....	6
Unguenti zinci oxidi.....	q. s. ad 100.

## 9.

Naphtolis.....	2
Sulphuris sublimati.....	4
Unguenti .....	q. s. ad 100.

(Kaposi.)

## 10.

Hydrastininae hydrochloratis.....	1
Zinci carbonatis.....	10
Unguenti zinci oxidi.....	q. s. ad 100.

## ALOPECIA AREATA.

## 1.

Acidi salicylici.....	3
Sulphuris præcipitati.....	15
Petrolati mollis.....	45
Adipis.....	q. s. ad 100.

Apply at night. Shampoo the head the next morning with a salicylated soap and apply friction with a soft brush, wet with the following:—

Hydrargyri chloridi corrosivi.....	1
Spiritus rosmarini.....	250
Alcoholis.....	q. s. ad 500.

Once a week the patches may be painted with:—

Spiritus gaultheriæ.....	50
Ætheris.....	q. s. ad 100.

(Feulard.)

## 2.

Acidi tannici ..	2
Olei ricini.....	50
Olei thymi.....	q. s.
Alcoholis absoluti.....	q. s. ad 100.

To be used as a head wash every evening. Once a week substitute the following:—

Acidi salicylici.....	1
Acidi acetici glacialis.....	50
Olei geranii.....	q. s.
Olei ricini.....	q. s. ad 100.

In addition to the above the head is to be washed once a week with soft soap.

(Rochon.)

3.

Tincturæ benzoini.....	1
Acidi salicylici.....	2
Alcoholis.....	q. s. ad 100.

Apply once or twice daily.

*(Kaposi.)*

4.

Olei cinnamomi.....	33
Ætheris.....	q. s. ad 100.

*(Gaucher.)*

5.

Spiritus ammoniæ.....	15
Extracti pilocarpi fluidi .....	30
Linimenti saponis mollis.....	q. s. ad 100.

*(Shoemaker.)*

6.

Tincturæ capsici.....	1
Balsami Peruviani.....	2
Alcoholis.....	q. s. ad 100.

*(Kaposi.)*

7.

Tincturæ cantharidis .....	10
Tincturæ capsici .....	10
Olei ricini ..	10
Alcoholis diluti.....	q. s. ad 100.

*(Ashby & Wright.)*

8.

Acidi acetici.....	10
Olei ricini .....	50
Alcoholis .....	q. s. ad 100.

9.

Acidi acetici glacialis.....	3
Chloralis.....	4
Ætheris .....	q. s. ad 100.

*(Gaucher.)*

## 10.

Quininæ.....	1
Acidi tannici.....	2
Olei resedæ.....	q. s.
Olei aurantii florum.....	q. s.
Unguenti aquæ rosæ.....	q. s. ad 100.

To be used occasionally if hair is dry.

## 11.

Tincturæ benzoini.....	2
Hydrargyri ammoniati.....	5
Olei rosæ.....	q. s.
Unguenti aquæ rosæ.....	q. s. ad 100.

Pomade. To be used if hair is very dry.

(*Kaposi.*)

## 12.

Chrysarobini.....	10
Ceræ albæ.....	30
Adipis lanæ hydrosi.....	q. s. ad 100.

For dispensary practice.

## 13.

Acidi salicylici.....	20
Ceræ albæ.....	25
Adipis lanæ hydrosi.....	q. s. ad 100.

For dispensary practice.

(*Unna.*)

## BITES AND STINGS OF INSECTS.

## 1.

Ammonii chloridi.....	10
Alcoholis.....	10
Aquæ... ..	q. s. ad 100.

## 2.

Hydrargyri chloridi corrosivi.....	1
Aquæ.....	q. s. ad 1000.



## 3.

Naphtolis..... 3  
 Aquæ.....q. s. ad 100.

## 4.

Potassii permanganatis..... 1  
 Aquæ.....q. s. ad 100.

## 5.

Aquæ hydrogenii dioxidi.....q. s.  
 For local application.

## 6.

Aquæ ammoniæ fortioris.....q. s.  
 Apply locally.

## 7.

Acidi borici..... 10  
 Unguenti aquæ rosæ.....q. s. ad 100.

## BURNS.

## 1.

## ANTISEPTIC CAGE TREATMENT.

A cradle or raised surface is placed so that an antiseptic dressing covers the wound but does not come in contact with it. The wound is dressed by lifting the dressing and washing the surface with a weak antiseptic solution.

(Benson.)

## 2.

## LONDON HOSPITAL TREATMENT.

A severely burned child is immediately wrapped in a blanket, without having its clothes removed. The bed is placed near a fire; hot-water bottles are applied; brandy and the tincture of opium are given. As the patient reacts, one part after another

is dressed with zinc ointment and covered with cotton. After the dressing the part is recovered with a blanket. Suppuration commences in about four days, and the parts are then redressed. It is at this period that diarrhœa is to be feared and should be treated at its inception. Never give aperients, but use enemata if constipation arises. In the interim, before the suppuration has commenced, the child is given two to three minims of the tincture of opium every four hours and one-half drachm of brandy every hour. The rule of never waking a child for its medicine is sufficient safeguard against an overdose. For a child two or three years old the first dose of the tincture should not be less than four or five minims, with two drachms of brandy, to be followed by three minims, if the child is crying, at the end of one hour.

*(Fenwick & Barnard.)*

3.

WATER-BED TREATMENT.

The patient is kept continually in Hebra's water-bed at a temperature agreeable to himself, and the wounds are treated surgically. The water is to be renewed two or three times a day. This bath can be imitated in private practice with a long bath tub, horsehair pillows, and blankets.

*(Hebra.)*

4.

TREATMENT OF POWDER BURNS.

Suppuration occurs in a few days. Apply wet compress for one-half of an hour or longer. A sharp razor is then passed over the surface, which cuts off the soft elevations containing the powder. Wash off with a stiff sponge moistened in a weak antiseptic solution. Clean and rewash until all coloring matter is removed.

*(Coombs.)*

## 5.

Potassii nitratis.....q. s.

Add a few teaspoonfuls to a basinful of water. The part affected should be bathed for from two to three hours. If preferred, compresses thoroughly wet with a saturated solution may be applied. The analgesic effect produced is due to the refrigerant action of the salt. If the pain returns when the temperature of the water is raised, more of the potash should be added or the compresses renewed.

(Poggi.)

## 6.

Sodii chloridi..... 1  
Aquæ.....q. s. ad 100.

Apply on compresses thoroughly wet with the solution. A window may be cut in the dressing through which the part may be kept constantly moist.

(Keen.)

## 7.

Bismuthi subnitratis.....q. s.  
Aquæ fervidæ.....q. s.

Mix to the consistency of a paste. Apply with a soft brush over the affected area. The bullæ in burns of the second degree are drained with antiseptic silk, and the raised epidermis is allowed to remain *in situ*. The original covering is added to from time to time, as it cracks or falls off, but no other dressing is used. Magnesia may be used in the same way.

(Osthoff.)

## 8.

Ichthyolis..... 6  
Zinci oxidi..... 30  
Cretæ præparatæ.....q. s. ad 100.

Apply once daily.

(Leistikow.)

## 9.

Acidi picrici.....	3
Aquæ.....	q. s. ad 200.

Immerse the affected part for five minutes and wrap in cotton, protecting it, if excoriated, with iodoform gauze. The stain can be removed with a solution of boric acid.

(*Thierry.*)

## 10.

Tincturæ opii.....	6
Liquoris plumbi subacetatis.....	6
Aquæ hamamelidis.....	12
Aquæ.....	q. s. ad 100.

## 11.

Sodii bicarbonatis.....	3
Aquæ.....	q. s. ad 100.

## 12.

Icthyolis.....	5
Zinci oxidi.....	10
Cretæ præparatæ.....	20
Amyli.....	20
Liquoris calcis.....	20
Olei lini.....	q. s. ad 100.

Apply once daily.

(*Leistikow.*)

## 13.

Aristolis.....	5
Olei olivæ.....	20
Sol. ad.	
Petrolati mollis.....	40
Adipis lanæ hydrosi.....	q. s. ad 100.

Do not apply until secretion ceases.

(*Haas.*)

## 14.

Cocainæ.....	1
Sodii bicarbonatis.....	5
Adipis lanæ hydrosi.....	15
Olei olivæ.....q. s. ad 100.	

(Shoemaker.)

## 15.

Argenti nitratis.....	50
Aquæ destillatæ.....q. s. ad 100.	

Nitrate of silver stick or this solution may be used to cauterize the exuberant granulations. The surface must first be moistened with a two per cent solution of cocaine.

---

### CHILBLAINS.

## 1.

Argenti nitratis.....	5
Aquæ.....q. s. ad 100.	

Use when itching is severe.

## 2.

Calcii chloridi.....	2
Aquæ.....q. s. ad 100.	

Use when itching is severe.

(Jacobi.)

## 3.

Talci.....	8
Resorcini.....	15
Mucilaginis acaciæ.....	40
Aquæ.....q. s. ad 100.	

Paint the surface.

(Boeck.)



4.

Iodi.....	1
Potassii iodidi.....	3
Glycerini.....q. s. ad 100.	

Paint the surface.

5.

Tincturæ iodidi.....	1
Olei ricini.....	5
Collodii.....q. s. ad 100.	

Paint the surface.

6.

Plumbi acetatis.....	12
Unguenti aquæ rosæ.....q. s. ad 100.	

*(Kaposi.)*

7.

Ichthyolis.....	25
Adipis lanæ hydrosi.....q. s. ad 100.	

For older children.

*(Hernane.)*

8.

Capsici.....	6
Olei amygdalæ expressi.....	24
Adipis lanæ hydrosi.....q. s. ad 100.	

Rub in with a piece of flannel.

*(Ashby & Wright.)*

## DERMATITIS GANGRÆNOSA.

1.

Hydrargyri chloridi corrosivi.....	1
Aquæ.....q. s. ad 500.	

2.

Chloralis.....	1
Aquæ.....q. s. ad 100.	

3.

Acidi salicylici.....	2
Aquæ.....	.q. s. ad 100.

4.

Acidi borici.....	20
Talci.....	40
Amyli.....	.q. s. ad 100.

5.

Acidi tannici.....	5
Petrolati mollis.....	.q. s. ad 100.

6.

Salolis .....	4
Petrolati mollis.....	.q. s. ad 100.

7.

Dermatolis.....	4
Petrolati mollis .....	.q. s. ad 100.

*(Gaucher.)*

## DERMATITIS EXFOLIATIVA NEONATORUM.

Hygienic measures, breast milk, and tonics constitute the general treatment. Protect the desquamating surface with cotton and by the application of oils and fats. An incubator should be used, if possible.

*(Ballantyne.)*

## ACUTE ECZEMA.

1.

Iridis.....	5
Zinci oxidi.. ..	5
Alumini.....	20
Amyli.....	.q. s. ad 100

## 2.

Plumbi carbonatis.....	10
Zinci oxidi.....	20
Bismuthi subnitratis.....	20
Talci.....	20
Amyli.....	q. s. ad 100.

Apply with a powder puff to the exposed parts of the skin, and to the intertriginous folds on pledgets of cotton thickly covered with powder. Change the pledgets as soon as they become warm and moist.

(*Kaposi.*)

## 3.

Zinci carbonatis.....	25
Plumbi carbonatis.....	25
Sulphuris sublimati.....	25
Marantæ pulveris.....	q. s. ad 100.

## 4.

Amyli.....	30
Plumbi carbonatis.....	35
Lycopodii.....	q. s. ad 100.

## 5.

Acidi tannici.....	1
Bismuthi subnitratis.....	8
Aquæ rosæ fortioris.....	q. s. ad 200.

(*Shoemaker.*)

## 6.

Plumbi acetatis.....	1
Alumini.....	5
Aquæ destillatæ.....	q. s. ad 100.

Dilute five times with water and apply on compresses.

## 7.

Acidi carbolici.....	1
Spiritus lavandulæ.....	20
Spiritus odorati.....	30
Alcoholis.....	q. s. ad 200.

Apply to the surface in case the pruritus is very severe. Follow immediately by the application of an inert powder. If there is not the slightest particle of moisture, tinctura rusci acts better. Apply in a very thin layer and follow with powder.

(Kaposi.)

## 8.

Zinci oxidi.....	10
Glycerini.....	20
Gelatini.....	25
Aquæ destillatæ.....	q. s. ad 100.

Cut off the quantity desired. Melt in a warm-water bath and apply with a brush. Only to be used when no moisture is present.

(Kaposi.)

## 9.

Naphtolis.....	1
Camphoræ.....	1
Cocainæ hydrochloratis.....	1
Zinci carbonatis.....	6
Unguenti aquæ rosæ.....	q. s. ad 100.

Use in acute infantile eczema.

## 10.

Creolini.....	2
Acidi borici.....	10
Zinci carbonatis.....	20
Marantæ.....	20
Adipis lanæ hydrosi.....	q. s. ad 100.

Use in acute infantile eczema.

(Shoemaker.)

## 11.

Dermatolis.....	6
Amyli.....	15
Petrolati mollis .....	q. s. ad 100.

Use in the squamous stage.

## 12.

Zinci oxidi .....	20
Amyli.....	20
Adipis .....	q. s. ad 100.

Use in the squamous stage.

## 13.

Hydrargyri ammoniati.....	10
Glyceriti amyli.....	q. s. ad 100.

Use in the squamous stage.

## 14.

Acidi tannici.....	5
Adipis lanæ hydroxi.....	30
Petrolati mollis .....	q. s. ad 100.

Use in the squamous stage if disease threatens to become chronic.

## 15.

Olei cadini.....	10
Petrolati mollis .....	q. s. ad 100.

Use in the squamous stage if disease threatens to become chronic.

## 16.

Acidi salicylici .....	2
Amyli.....	8
Zinci oxidi.....	10
Adipis lanæ hydroxi.....	15
Petrolati mollis .....	q. s. ad 150.

Use in the squamous stage if disease threatens to become chronic.

(Gaucher.)



## CHRONIC ECZEMA.

1.

Liquoris potassii arsenitis.....	3
Syrupi aurantii florum.....	25
Olei morrhuæ.....	50
Mucilaginis.....	q. s.
Aquæ.....	q. s. ad 100.

One teaspoonful three times a day, after meals, for a child less than one year old.

(*Ashby & Wright.*)

2.

Syrupi ferri iodidi.....	20
Extracti malti fluidi.....	q. s. ad 100.

One-half teaspoonful to two teaspoonfuls three times a day.

(*Shoemaker.*)

3.

Plumbi acetatis.....	2
Aquæ.....	q. s. ad 100.

Apply as a lotion.

4.

Alumini.....	3
Aquæ.....	q. s. ad 100.

Apply as a lotion.

5.

Argenti nitratis.....	5
Aquæ.....	q. s. ad 100.

Apply as a lotion.

6.

Acidi tannici.....	4
Glycerini.....	50
Aquæ.....	q. s. ad 100.

Apply as a lotion.

## 7.

Chloralis.....	3
Glycerini.....	50
Aquæ.....	q. s. ad 100.

Apply as a lotion when the area affected is small.

(*Gaucher.*)

## 8.

Unguenti cum styrace.....	20
Olei olivæ.....	q. s. ad 100.

(*Vidal.*)

## 9.

Zinci oxidi.....	50
Adipis lanæ hydrosi.....	q. s. ad 100.

In eczema universalis the child is covered from head to foot with soft cotton compresses thickly spread with the ointment. He is held in bed by broad straps drawn across his legs, abdomen, chest, and shoulders, thus binding his arms to his side and keeping his legs in extension. Soft, heavily padded sand bags are placed one on either side of the head, to prevent any movement and consequent irritation. A nurse is in continual attendance at first, and soothing drugs may be given if necessary.

(*Rotch.*)

## 10.

Cretæ præparatæ.....	3
Sulphuris sublimati.....	15
Olei fagi.....	15
Saponis mollis.....	30
Adipis.....	q. s. ad 100.

Apply thinly with a brush.

(*Hebra.*)

## 11.

Olei rusci.....	1
Zinci oxidi.....	5
Petrolati mollis.....	50
Adipis lanæ hydrosi.....	q. s. ad 100.

(*Baginsky.*)

## 12.

Hydrargyri ammoniati.....	3
Balsami Peruviani.....	15
Unguenti zinci oxidi.....	q. s. ad 100.

(Saalfeld.)

## 13.

Olei cadini.....	25
Petrolati mollis.....	25
Adipis lanæ hydrosi.....	q. s. ad 100.

The oil of cade should be pure and made from the wood.  
The coal-tar product is much inferior in its effect.

## 14.

Ichthyolis.....	10
Petrolati mollis.....	q. s. ad 100.

## 15.

Naphtolis....	2
Petrolati mollis.....	q. s. ad 100.

To be used with great care in treating very young children.

## 16.

Acidi borici.....	8
Hydrargyri ammoniati.....	8
Glyceriti amyli.....	q. s. ad 100.

Use in eczema impetigo.

## 17.

Hydrargyri oxidi flavi.....	3
Petrolati mollis.....	q. s. ad 100.

Use in eczema impetigo.

(Gaucher.)

ECZEMA ANI.

1.

Chloralis.....	3
Glycerini.....	50
Aquæ.....	q. s. ad 100.

(Gaucher.)

2.

Acidi tannici.....	4
Adipis.....	q. s. ad 100.

3.

Extracti krameriæ.....	12
Petrolati mollis.....	q. s. ad 100.

4.

Mentholis.....	1
Petrolati mollis.....	q. s. ad 100.

To be used if pruritus is severe.

5.

Cocainæ hydrochloratis.....	1
Unguenti.....	100.

---

ECZEMA AURIUM.

1.

Sodii carbonatis.....	2
Glycerini.....	q. s. ad 100.

Pour into the external auditory meatus to loosen the scales, and remove them with a syringe of warm water.

2.

Zinci sulphatis.....	1
Glycerini.....	q. s. ad 200.

Warm and pour into the external auditory meatus, using from ten to fifteen drops daily.

3.

Argenti nitratis.....	1
Aquæ destillatæ .....	q. s. ad 100.

Paint the surfaces with a brush. Cease the application if any pain arises.

(Gruber.)

4.

Unguenti diachyli.....	50
Petrolati mollis .....	q. s. ad 100.

Apply on cloths.

5.

Glycerini.....	10
Olei fagi .....	20
Unguenti aquæ rosæ .....	q. s. ad 100.

Apply in chronic cases when the epidermis is thickened.

(Politzer.)

6.

Creolini.....	1
Aristolis.....	4
Unguenti hydrargyri nitratis.....	35
Unguenti zinci oxidi.....	q. s. ad 100.

(Shoemaker.)

## ECZEMA CAPITIS.

1.

Naphtolis.....	1
Olei olivæ.....	q. s. ad 100.

Use to soften the crusts.

(Kaposi.)

2.

Acidi salicylici .....	1
Olei morrhuæ.....	q. s. ad 100.

Apply locally to soften the crusts. Wash off the softened masses every day or every other day with the liniment of soft soap.

(Baginsky.)



## 3.

Acidi carbolicī.....	1
Balsami Peruviani.....	2
Olei olivæ.....q. s. ad 100.	

Use to soften the crusts. If the inflammation is severe cold douches or compresses may be used.

*Kaposi.*)

## 4.

Zinci oxidi.....	5
Petrolati mollis.....	50
Adipis lanæ hydrosi .....q. s. ad 100.	

Allow to remain on the scalp for one or two days, then apply Lassar's paste. When desired, remove the paste and crusts with a two per cent boric acid ointment in vaseline.

*(Baginsky.)*

## 5.

Acidi salicylici.....	2
Zinci oxidi.....	25
Amyli .....	25
Petrolati mollis.....q. s. ad 100.	

Apply in a thick layer.

*(Lassar.)*

## 6.

Zinci oxidi.....	25
Amyli.....	25
Petrolati mollis.....q. s. ad 100.	

Apply in a thick layer. An inert powder may be applied over the paste.

*(Lassar.)*

## 7.

Sulphuris præcipitati .....	4
Zinci oxidi.....	25
Amyli .....	25
Adipis lanæ hydrosi.....	25
Petrolati mollis.....q. s. ad 100.	

Apply in a thick layer.

*(Ihle.)*

## ECZEMA LABIORUM.

1.

Hydrargyri chloridi corrosivi..... 1  
 Tincturæ benzoini.....q. s. ad 500.  
 (*Taylor.*)

2.

Acidi salicylici..... 2  
 Petrolati mollis.....q. s. ad 100.  
 (*Kaposi.*)

3.

Hydrargyri oxidi flavi..... 3  
 Petrolati mollis... ..q. s. ad 100.  
 (*Gaucher.*)

4.

Thymolis..... 1  
 Unguenti aquæ rosæ.....q. s. ad 100.

## ECZEMA NARIUM.

1.

Acidi salicylici..... 1  
 Olei amygdalæ expressi.....q. s. ad 1000.  
 Introduce into nasal cavities on tampons of cotton.  
 (*Besnier.*)

2.

Glyceriti plumbi subacetatis..... 50  
 Unguenti aquæ rosæ.....q. s. ad 100.  
 (*Hardaway.*)

3.

Zinci oxidi..... 10  
 Olei theobromatis.....q. s. ad 100.

Insert in nasal cavity. Each suppository should contain one grain of the oxide.

(*Neumann.*)

## ECZEMA PALPEBRARUM.

1.

Acidi borici.....	5
Glycerini.....	5
Unguenti... ..	.q. s. ad 100.

2.

Hydrargyri oxidi flavi.....	1
Unguenti aquæ rosæ.....	.q. s. ad 100.

*(Kaposi.)*

3.

Aristolis.....	2
Unguenti aquæ rosæ.....	.q. s. ad 100.

4.

Europheni.....	3
Unguenti.....	.q. s. ad 100.

*(Shoemaker.)*

## ECZEMA UMBILICI.

1.

Thymolis.....	1
Cinchonæ rubræ pulveris.....	50
Bismuthi subnitratis.....	.q. s. ad 100.

*(Shoemaker.)*

2.

Liquoris plumbi subacetatis.....	50
Aquæ.....	.q. s. ad 100.

Apply on tampons moistened with the solution.

3.

Cocainæ hydrochloratis.....	1
Extracti belladonnæ foliorum alcoholici.....	4
Acidi tannici.....	6
Unguenti zinci oxidi.....	.q. s. ad 100.

**ERYSIPELAS.**

1.

Quininæ sulphatis .....	1
Acidi sulphurici, q. s. ad sol.	
Syrupi. ....	50
Aquæ destillatæ.....q. s. ad 100.	

A teaspoonful every two hours for a nursing child. If preferred, the tincture of iron may be given in doses of from five to ten drops every two hours, and to nursing children two drops every two hours.

2.

Pilocarpinæ hydrochloratis.....q. s.	
--------------------------------------	--

Use subcutaneously in doses of from one-tenth of a grain to three-tenths of a grain each. Repeat after two, eight, and ten hours. It may also be given internally at the same time.

*(Bau.)*

3.

Sodii benzoatis.....q. s.	
---------------------------	--

Give from five to ten grains, in divided doses, during twenty-four hours.

*(Schiller.)*

4.

**ANTISTREPTOCOCCIC SERUM TREATMENT.**

Give in hypodermatic doses of ten cubic centimetres each. These doses are of a serum prepared by Marmorek which is constant in strength and character.

5.

Ichthyolis. ....	50
Adipis lanæ hydrosi.....q. s. ad 100.	

Employ multiple scarification of the affected area, reaching into the corium and partly into the subcutaneous tissue. The surface is thoroughly washed with an antiseptic solution, and as much of the œdematous fluid as possible is expressed. An incision reaching to the corium is made, completely encircling the affected area. The ointment is now rubbed in and applied plentifully over the surface, which is covered with gauze and

a thick layer of cotton. The treatment is continued for a few days after the temperature has subsided. In most cases anæsthesia is necessary.

(*Felsenthal.*)

## 6.

Alcoholis absoluti..... q. s.

Apply continuously on thoroughly wet compresses or cotton covered with some impermeable material. Change every fifteen or twenty minutes, and apply lanoline when scaling occurs.

(*Sarysdorff.*)

## 7.

Ichthyolis..... 25

Liquoris gutta-perchæ.....q. s. ad 100.

Apply with a brush. Application should extend beyond the affected area by three-quarters of an inch. If the face is affected, surround the ears, even if not affected, with a band of the application.

## 8.

Extracti ergotæ ..... 1

Extracti ergotæ fluidi ..... 10

Adipis lanæ hydrosi ..... 20

Adipis.....q. s. ad 100.

Apply locally in cases of facial erysipelas.

(*Lees.*)

## ERYTHEMA.

## 1.

Zinci oxidi ..... 10

Talci..... 50

Magnesiæ.....q. s. ad 100.

(*Baginsky.*)

## 2.

Acidi salicylici..... 1

Magnesiæ..... 50

Amyli.....q. s. ad 100.



3.

Acidi borici .....	1
Magnesiæ .....	50
Amyli.....	.q. s. ad 100.

4.

Zinci oxidi .....	3
Calaminæ præparatæ.....	3
Liquoris calcis.....	.q. s. ad 100.

*(Rotch.)*

5.

Acidi borici .....	4
Glycerini.....	30
Aquæ rosæ fortioris .....	.q. s. ad 100.

*(Shoemaker.)*

## ERYTHEMA NODOSUM.

1.

Sodii salicylatis.....	10
Amyli.....	.q. s. ad 100.

Apply on wet cloths. Quinine or salicylate of soda may be given internally.

*(Monti.)*

2.

Argenti nitratis.....	6
Aquæ.....	.q. s. ad 100.

*(Jacobi.)*

3.

Olei gaultheriæ.....	.q. s.
----------------------	--------

Use as an inunction.

4.

Potassii iodidi.....	25
Glycerini sive	
Adipis lanæ hydrosi.....	.q. s. ad 100.

Apply locally. If preferred, lead water (pure or combined with opium) may be applied.

## FAVUS.

## 1.

Acidi carbolicæ.....	3
Aquæ.....	q. s. ad 100.

The head is washed and shaved, and epilation is performed. Compresses wet with this solution are then applied continuously from morning until evening, but are removed at night. A coil is laid on the compresses, through which water passes at a temperature of from 125° to 130° F. The compresses consist of about sixteen thicknesses of gauze. If any signs of carbolic poisoning arise, change the solution to a one-quarter per cent sublimate solution, or the latter may be used from the first. Treatment is continued during from eight to fourteen days, when boric vaseline is applied. The child can move about in bed, but the coil must cover the entire scalp, and there should be as little space as possible between the tubes of the coil.

(Zinsser.)

## 2.

Acidi borici.....	3
Alcoholis.....	50
Ætheris.....	q. s. ad 100.

(Shoemaker.)

## 3.

Thymolis.....	2
Chloroformi.....	20
Olei olivæ.....	q. s. ad 100.

Apply on compresses three times daily. When microscopic examination of the scales shows that there are no more fungi present, paint the surface daily during eight days with a thirty per cent solution of glycerine and the tincture of iodine.

(Gouladze.)

## 4.

Losophani.....	10
Unguenti aquæ rosæ.....	20
Adipis lanæ hydrosi.....	q. s. ad 100.

Apply twice daily.

(Saalfeld.)

## 5.

Potassii carbonatis.....	2
Sulphuris sublimati.....	8
Tincturæ iodidi.....	25
Olei picis liquidæ.....	25
Adipis.....	q. s. ad 100.

Apply once daily. Continue until active desquamation is established, and allay inflammation with a paste of zinc oxide, starch, and salicylic acid.

(*Tsitrine.*)

## FRECKLES AND OTHER PIGMENTATIONS.

## 1.

Hydrargyri chloridi corrosivi.....	1
Aquæ.....	q. s. ad 100.

Apply on closely fitting cloths. The patient is to remain in the horizontal position and the cloths are to be kept moist for four hours. Pierce the bullæ at their lower edges, so as to cause them to collapse, and apply an inert powder. The crust will fall off within a week, and the newly-formed epidermis will be unpigmented. This method is very painful and should not be used with children under ten years of age in cases of freckles on the face.

(*Kaposi.*)

## 2.

Glycerini.....	2
Naphtolis.....	4
Linimenti saponis mollis.....	q. s. ad 100.

Apply once daily as a wash.

## 3.

Hydrargyri chloridi corrosivi.....	1
Tincturæ benzoini.....	90
Aquæ aurantii florum fortioris.....	900
Emulsi amygdalæ.....	3000.

Apply once daily as a wash.

4.

Saponis mollis.....q. s.

Wash the face daily with the soft soap and water, and then paint it with either a weak salt or acetic acid solution.

5.

Hydrargyri ammoniati..... 15

Bismuthi subnitratiss..... 15

Glyceriti amyli.....q. s. ad 100.

Apply every other day.

(Hager.)

6.

Acidi salicylici..... 2

Tincturæ benzoini..... 4

Unguenti aquæ rosæ.....q. s. ad 100.

Spread on gauze and leave on over night in cases of freckles.

7.

Bismuthi subcarbonatis..... 10

Talcis..... 25

Sulphuris præcipitati..... 30

Olei rosæ.....q. s.

Barii dioxidi.....q. s. ad 100.

To be used if the skin becomes red and scaly in the course of healing.

(Kaposi.)

---

## FURUNCLE.

1.

Extracti colchici radicis.....q. s.

Give in doses of from one-third to one-half of a grain two or three times daily. Locally, the area affected is well washed with spirits of camphor, and Vidal's red plaster is then applied.

(Brocq.)

## 2.

## GALVANO-CAUTERIZATION TREATMENT.

Use a thin platinum wire point about one-quarter of an inch in length and one-twenty fifth of an inch in diameter, at white heat. It is passed into the follicle to the root of the hair. If applied early enough it is abortive in its action. If used later, when a little pus is already present, the application must be prolonged until the pus is entirely dried up by the heated wire. A thermo-cautery may be used, but it is more painful and causes greater destruction of tissue.

(*Loewenburg.*)

## 3.

Camphoræ ..... 55  
Acidi carbolici ..... q. s. ad 100.

Wash well with alcohol and apply a tampon moistened with this solution. Renew every twenty-four hours until active inflammation has subsided, then substitute a four-fifths per cent ointment of the yellow oxide of mercury. If an incision is necessary, incise to periosteum and reapply tampon. Care should be taken in cases of small children, on account of the amount of carbolic acid employed. To be used for small furuncles in the ear only.

(*McCuen Smith.*)

## 4.

Acidi salicylici..... 10  
Emplastri saponis..... q. s. ad 100.

Apply to the affected parts. A two per cent solution of carbolic acid may be used first.

(*Newberger.*)

## HYPERIDROSIS.

## 1.

Agaracini ..... q. s.

Give one-thirtieth of a grain every three hours.



2.

Pulveris agarici albi.....q. s.

Give from one to two grains three times daily.

3.

Atropinæ sulphatis.....q. s.

Give in doses of from one-three-hundredth of a grain to one-one-hundred-and-fiftieth of a grain twice daily.

4.

Zinci oxidi..... 10

Amyli.....q. s. ad 100.

Apply on lint in the interspaces between the fingers, toes, genital folds, and axillæ. This powder may also be placed in the stockings.

5.

Naphtolis ..... 1

Acidi salicylici..... 2

Amyli.....q. s. ad 100.

Apply as in No. 4.

*(Kaposi.)*

6.

Acidi tannici..... 1

Alcoholis.....q. s. ad 200.

Use as a local wash in cases of hyperidrosis of axillæ, genitals, palms of the hands, or soles of the feet.

7.

Hydrargyri chloridi corrosivi..... 1

Aquæ.....q. s. ad 800

Use as a local wash.

8.

Naphtolis ..... 1

Spiritus odorati. . . . . 3

Alcoholis.....q. s. ad 300.

Apply locally as a wash.

## 9.

Balsami Peruviani . . . . .	1
Acidi formici . . . . .	5
Chloralis . . . . .	5
Alcoholis . . . . .	q. s. ad 100.

Apply with cotton that can be destroyed subsequently. In cases of slight severity the chloral may be omitted. If a more powerful action is desired, add one per cent of trichloracetic acid. This solution may also be used in the form of a spray in cases of general hyperidrosis.

(Huesner.)

## 10.

Unguenti picis liquidæ . . . . .	50
Unguenti sulphuris . . . . .	q. s. ad 100.

Wash the parts with juniper or tar soap. Apply on cloths.

(Van Harlingen.)

## 11.

Emplastri plumbi . . . . .	20
Olei olivæ . . . . .	q. s. ad 100.
Sub leni igni et addendo pauxill. aqu. font. coque ut f. ungu. consistent. spissior.	
Adde: Olei lavandulæ florum . . . . .	2

Apply once daily on cloths. Dry the feet with powder or lint, but do not wash them. Repeat for from ten to fourteen days. Bathe when exfoliation occurs. Use in severe cases only.

(Hebra.)

## ICHTHYOSIS.

## 1.

Acidi salicylici . . . . .	2
Adipis lanæ hydrosi . . . . .	50
Petrolati mollis . . . . .	q. s. ad 100.

2.

Acidi carbolici.....	1
Adipis .....	q. s. ad 100.

3.

Resorcini.....	4
Petrolati mollis.....	50
Adipis lanæ hydrosi.....	q. s. ad 100.

*(Gaucher.)*

4.

Tincturæ iodidi.....	25
Glycerini.....	40
Olei theobromatis.....	q. s. ad 100.

*(Perez.)*

5.

Acidi tartarici.....	3
Glyceriti amyli.....	q. s. ad 100.

Apply at first every evening, later every two or three days, then once a week, and finally from time to time. When this is not applied, use the glycerite of starch every evening.

*(Vidal.)*

6.

Naphtolis.....	2
Petrolati mollis.....	q. s. ad 100.

Rub in sparingly once or twice daily. Naphthol soap may also be used every other day. In young children inunctions of pure soft soap may be used exclusively.

*(Kaposi.)*

## IMPETIGO.

1.

Hydrargyri chloridi corrosivi.....	1
Aquæ.....	q. s. ad 100.

Remove the crusts. Stop all hemorrhage with tampons. Rub the affected areas, especially the borders of the pustules, with this solution. Repeat until no new pustules occur. In most cases two to three daily successive applications are sufficient. If no new pustules appear, those already treated heal rapidly under a zinc ointment or a plaster.

(Neebe.)

2.

Olei cadini .....	50
Olei olivæ.....	q. s. ad 100.

3.

Hydrargyri oxidi flavi.....	5
Petrolati mollis .....	q. s. ad 100.

4.

Salolis.....	10
Petrolati mollis .....	q. s. ad 100.

5.

Acidi borici .....	8
Zinci oxidi .....	12
Glyceriti amyli.....	q. s. ad 100.

6.

Acidi borici .....	8
Hydrargyri ammoniati .....	8
Petrolati mollis .....	q. s. ad 100.

7.

Acidi salicylici .....	1
Zinci oxidi .....	10
Petrolati mollis .....	q. s. ad 100.

(Gaucher.)

## INTERTRIGO.

1.

Acidi chromici .....	3
Aquæ.....	q. s. ad 100.

Wash and dry the parts carefully. Paint with this solution, and cover it with some bland powder. Repeat every three or four days. Not more than two or three applications are required generally. If there is great irritation of the parts, some weak antiseptic solution should be used first. This treatment prevents recurrence, by tanning the skin.

(Brault.)

2.

Argenti nitratis .....	2
Aquæ.....	q. s. ad 100.

To be used in very obstinate cases limited to the anus or genitals. A three-quarters per cent solution of bichloride of mercury may be used if preferred. Bichloride or bran baths at a temperature of 80° F. are often effective.

(Hench.)

3.

Ichthyolis .....	1
Aquæ.....	q. s. ad 200.

Apply locally if the parts are infected. Powder is to be applied over this. The parts should be kept separated with cotton covered with starch.

4.

Resorçini.....	1
Aquæ... ..	q. s. ad 100.

Apply locally as in No. 3.

5.

Olei olivæ.....	50
Liquoris calcis.....	q. s. ad 100.



## 6.

Bismuthi subnitrat. ....	1
Adipis lanæ hydrosi. ....	q. s. ad 100.

## 7.

Acidi salicylici. ....	1
Cetacei. ....	10
Adipis lanæ hydrosi. ....	q. s. ad 100.

(Widerhofer.)

## 8.

Ichthyolis. ....	2
Olei olivæ. ....	20
Adipis lanæ hydrosi. ....	q. s. ad 100.

(Monti.)

## 9.

Zinci oxidi. ....	10
Petrolati mollis. ....	q. s. ad 100.

(Neumann.)

## LUPUS.

## 1.

Argenti nitratis. ....	30
Aquæ destillatæ. ....	q. s. ad 100.

Paint the affected areas.

## 2.

Acidi lactici. ....	80
Aquæ. ....	q. s. ad 100.

## 3.

Acidi tartarici. ....	2
Alcoholis. ....	15
Acidi carbolic. ....	25
Olei ricini. ....	25
Camphoræ. ....	q. s. ad 100.

(Gaucher.)

## 4.

Iodi.....	25
Potassii iodidi .....	25
Glycerini.....	q. s. ad 100.

Paint the surface affected.

## 5.

Acidi arsenosi.....	3
Hydrargyri sulphidi rubri .....	12
Unguenti hydrargyri oxidi rubri.....	q. s. ad 100.

Spread on linen and cut into sizes desired.

(*Hebra.*)

## 6.

Hydrargyri chloridi corrosivi.....	1
Ætheris, q s. ad sol.	
Petrolati mollis .....	q. s. ad 100.

To be applied over small surfaces only.

(*Doutretpont.*)

## 7.

Hydrargyri iodidi rubri.....	50
Adipis.....	q. s. ad 100.

Apply locally. Warm before using.

(*Hardy.*)

## 8.

Acidi pyrogallici.....	10
Petrolati mollis.....	q. s. ad 100.

(*Schwimmer.*)

## 9.

Resorcini.....	35
Petrolati mollis.....	q. s. ad 100.

(*Bertarelli.*)

10.

Creosoti .....	4
Acidi salicylici.....	20
Emplastri plumbi.....q. s. ad 100.	

*(Besnier.)*

11.

Potassii chloridi.....	25
Potassii nitratis.....	25
Zinci chloridi .....	q. s. ad 100.

Melt together in the form of a pencil. Apply locally.

*(Koebner.)*

12.

Zinci chloridi.....	50
Aquæ .....	q. s.
Amyli.....	q. s. ad 100.

Make into a solid mass. At the time of application, cut off the amount desired.

*(Canquoin.)*


---

## LUPUS ERYTHEMATOSUS.

1.

Iodi ... ..	10
Potassii iodidi.....	20
Aquæ .....	q. s. ad 100.

Apply with a brush once or twice daily until sufficient irritation is induced.

*(Hardy.)*

2.

Ichthyolis.....	15
Petrolati mollis .....	q. s. ad 100.

Apply over night. Wash with warm water in the morning and apply some bland or antiseptic ointment, as oxide of zinc or a ten per cent boric acid ointment in vaseline. If irritation is then extreme, continue with the non-irritating ointment. Otherwise use the ichthyol for some nights.

3.

Acidi salicylici .....	3
Petrolati mollis .....	.q. s. ad 100.

Apply as in No. 2.

4.

Resorcini. ....	10
Petrolati mollis .....	.q. s. ad 100.

Apply as in No. 2.

(*Gaucher.*)

## MILIARIA.

1.

Camphoræ .....	6
Bismuthi subnitratis.....	15
Zinci oxidi.....	.q. s. ad 100.

2.

Calaminæ præparatæ .....	3
Zinci oxidi.....	6
Glycerini.....	12
Liquoris calcis.....	24
Aquæ rosæ fortioris.....	.q. s. ad 100.

Apply on a piece of muslin dipped in the solution. Bandage it on if necessary. Calomel or saline laxatives may also be given, or diuretics in the form of the nitrate of potash or the sweet spirits of nitre.

(*Holt.*)

3.

Liquoris plumbi subacetatis.....	1
Tincturæ opii.....	4
Aquæ.....	.q. s. ad 100.

Apply locally and follow with the application of a boric acid powder. To be used in cases of lichen strophulous.

(*Crocker.*)

## NÆVUS VASCULOSUS.

## 1.

## VACCINE-LYMPH TREATMENT.

Three to six superficial cross-incisions are made in the tumor, and the vaccine is introduced in the same manner as in the ordinary process of vaccination. This treatment is to be used in the case of a child who has never been vaccinated. It is likely to leave a larger scar than the electrolytic needle, but is often effective in cases of nævi of moderate size. The child is usually vaccinated in some other part of his body at the same time.

## 2.

Hydrargyri chloridi corrosivi ..... 10  
Collodii.....q. s. ad 100.

Apply with a brush to the area affected. This is recommended highly by various writers for the treatment of nævus flammeus and araneus, as well as nævus tuberosus, if the latter is small.

## DISEASES OF THE NAILS.

## ECZEMA OF THE NAILS.

## 1.

Acid salicylici..... 5  
Petrolati mollis.....q. s. ad 100.

## 2.

Ichthyolis..... 10  
Petrolati mollis .....q. s. ad 100.

## 3.

Unguenti cum styrace..... 30  
Olei olivæ.....q. s. ad 100.



4.

Naphtolis.....	4
Petrolati mollis.....	q. s. ad 100.

*(Gaucher.)*

5.

Oleati stanni.....	6
Petrolati mollis.....	q. s. ad 100.

*(Shoemaker.)*

## FAVUS OF THE NAILS.

1.

Hydrargyri chloridi corrosivi.....	3
Aquæ.....	q. s. ad 300.

Remove or scrape and file away the nail until perforated. Curette the small yellow areas. Apply the solution on small cotton tampons. Keep in position with a bandage.

2.

Naphtolis.....	5
Petrolati mollis.....	q. s. ad 100.

3.

Acidi pyrogallici.....	5
Adipis lanæ hydrosi.....	q. s. ad 100.

4.

Acidi chrysophanici.....	8
Adipis lanæ hydrosi.....	q. s. ad 100.

*(Gaucher.)*

## PSORIASIS OF THE NAILS.

1.

Gallanolis.....	10
Alcoholis.....	q. s. ad 100.

2.

Acidi chrysophanici..... 10  
 Chloroformi.....q. s. ad 100.

3.

Acidi pyrogallici..... 10  
 Ætheris.....q. s. ad 100.

4.

Gallanolis..... 15  
 Petrolati mollis.....q. s. ad 100.

5.

Gutta-Perchæ..... 10  
 Chloroformi.....q. s. ad 100.

Apply with a soft brush.

6.

Gutta-Perchæ..... 10  
 Gallanolis..... 10  
 Chloroformi.....q. s. ad 100.

Paint the affected area.

(*Gaucher.*)

## RINGWORM OF THE NAILS.

1.

Hydrargyri chloridi corrosivi..... 1  
 Alcoholis.....q. s. ad 200.

Scrape away as much of the nail as possible and apply as a wet compress once daily. In the intervals between the applications of this solution, apply a boric acid ointment.

(*Gaucher.*)

2.

Iodi..... 1  
 Potassii iodidi..... 2  
 Aquæ destillatæ.....q. s. ad 1000.

Apply constantly as a wet dressing, and keep in place with rubber finger stalls.

(*Sabouraud.*)

	3.	
Acidi pyrogallici.....		50
Olei olivæ.....	q. s. ad 100.	

Bandage in position. The nail will become movable in a few days, when apply an antiseptic dressing twice daily.

(*Pellizari.*)

## PHTHIRIASIS.

### PHTHIRIASIS CAPITIS.

1.

Linimenti saponis mollis.....q. s.

The hair is to be cut, if possible, and thoroughly washed with the liniment. When the hair cannot be cut, the head is to be thoroughly washed and a one-third per cent sublimate solution in vinegar is then to be applied. Spirits of camphor or warm vinegar may be used in the same way. After twenty-four hours the head is to be washed again with soft soap and thoroughly combed with a fine comb wet in vinegar.

(*Gaucher.*)

2.

Acidi carbolici ..... 2  
Olei olivæ.... q. s. ad 100.

Rub in well and cover the head with a cloth.

(*Monti.*)

3.

Balsami Peruviani..... 3  
Olei olivæ ..... 35  
Petrolei.....q. s. ad 100.

Rub the scalp thoroughly with this solution. Cover it with flannel over night and wash with soft soap in the morning. Repeat for two or three days. Nits are to be removed by washing the hair with vinegar and combing it with a fine comb.

4.

Naphtolis ..... 1  
 Olei olivæ.....q. s. ad 100.

(Kaposi.)

## PHTHIRIASIS CORPORIS.

1.

Acidi carbolici ..... 1  
 Aquæ..... q. s. ad 100.

The patient is to be well bathed; his clothes are to be exposed to a high temperature; and a complete change of clothing is to be made. This alone is often effective. In a chronic case with severe cutaneous lesions, fumigation, sulphur baths, a mild solution of sublimate, or the above-mentioned solution may be used. Jamieson recommends wearing a piece of roll sulphur, about the size of a pigeon's egg, next to the skin. It is to be wrapped up in gauze and worn day and night.

## PHTHIRIASIS PUBIS.

1.

Hydrargyri chloridi corrosivi ..... 1  
 Alcoholis.....q. s. ad 500.

Apply locally and wash the parts with warm water. One application is generally sufficient, but it may be necessary to repeat it two or three times.

2.

Hydrargyri chloridi corrosivi ..... 1  
 Aquæ destillatæ.....q. s. ad 100.

Apply as in No. 1.

3.

Hydrargyri ammoniati ..... 10  
 Unguenti aquæ rosæ.....q. s. ad 100.

(Kaposi.)

## PRURIGO.

## 1.

Acidi carbolici.....	1
Aquæ destillatæ.....	q. s. ad 100.

Inject one small hypodermatic syringeful every day or every other day. Use lanoline externally and warm baths at the same time.

(*Monti.*)

## 2.

Calcis .....	8
Sulphuris sublimati.....	16
Coq. c. aq. fervid.....	160
Ad reman.....	100.
Cola et filtra.	

To be well rubbed into the skin.

(*Henoch.*)

## 3.

Petrolei.....	50
Glycerini.....	q. s. ad 100.

For nursing children. Rub in with flannel wet with the solution. In private practice substitute the balsam of Peru for the petroleum and apply with a brush.

(*Monti.*)

## 4.

Naphtolis .....	1
Petrolati mollis .....	q. s. ad 100.

Apply two or three times daily. Rub dry with cotton after the last daily application. Bathe the patient after ten or twelve applications.

(*Baginsky.*)

## 5.

Unguenti diachylon.....	50
Petrolei.....	q. s. ad 100.

Apply on flannel three times daily if itching is severe.



## PRURITUS CUTANEUS.

1.

Camphoræ.....	2
Zinci oxidi.....	50
Amyli.....	q. s. ad 100.

2.

Hydrargyri chloridi corrosivi.....	1
Alcoholis.....	50
Aquæ laurocerasi.....	50
Aquæ.....	q. s. ad 1000.

3.

Chloralis.....	2
Glycerini.....	15
Alcoholis.....	15
Aquæ.....	q. s. ad 100.

4.

Mentholis.....	1
Glycerini.....	20
Alcoholis.....	q. s. ad 200.

*(Kaposi.)*

5.

Acidi carbolici.....	1
Zinci oxidi.....	15
Petrolati mollis.....	q. s. ad 100.

6.

Camphoræ.....	10
Petrolati mollis.....	q. s. ad 100.

7.

Cocainæ hydrochloratis.....	1
Petrolati mollis.....	q. s. ad 100.

*(Gaucher.)*

## PSORIASIS.

## 1.

Olei cadini.....	10
Glyceriti amyli.....	.q. s. ad 100.

## 2.

Acidi chrysophanici.....	5
Petrolati mollis.....	q. s. ad 100.

## 3.

Acidi salicylici.....	3
Olei cadini.....	5
Petrolati mollis.....	.q. s. ad 100.

## 4.

Sulphuris sublimati.....	12
Olei cadini.....	12
Petrolati mollis.....	.q. s. ad 100.

## 5.

Hydrargyri ammoniati.....	10
Petrolati mollis.....	.q. s. ad 100.

## 6.

Hydrargyri oxidi rubri.....	6
Petrolati mollis.....	.q. s. ad 100.

(Gaucher.)

## 7.

Acidi salicylici.....	2
Sulphuris præcipitati.....	10
Zinci oxidi.....	20
Amyli.....	20
Petrolati mollis.....	.q. s. ad 100.

To be used before the application of chrysarobin, combined with protracted baths.

(Lassar.)

8.

Gallacetophenoni.....	10
Unguenti.....	q. s. ad 100.

Apply as an ointment. Non-irritating. Effective, but slow in its action.

(Gall.)

9.

Chrysarobini.....	20
Zinci oxidi.....	20
Adipis lanæ hydrosi.....	30
Petrolati mollis.....	q. s. ad 100.

To be used on small areas only.

(Neisser.)

10.

Chrysarobini.....	8
Liquoris gutta-perchæ.....	8
Chloroformi.....	q. s. ad 100.

To be applied to affected parts twice a week.

(Ashby &amp; Wright.)

---

## PURPURA RHEUMATICA.

1.

Extracti ergotæ.....	q. s.
----------------------	-------

Give one-tenth of a grain in pill form three times a day. It may be given subcutaneously in doses of one and one-half grains every other day, but is very liable to cause abscesses.

2.

Extracti opii.....	4
Unguenti aquæ rosæ.....	q. s. ad 100.

Not to be applied over a large area in cases of young children.

3.

Extracti opii.....	6
Emplastri saponis.....	q. s. ad 100.

Apply on linen and bandage in place. Not to be used in cases of very young children.

## RINGWORM.

1.

Formalini.....	15
Alcoholis.....	q. s. ad 100.

Shave the hair around the margin of the patches and rub in the fluid vigorously with a brush for ten minutes every other day. Apply four times. Salter recommends a forty per cent solution.

2.

Acidi borici.....	3
Tincturæ camphoræ.....	3
Glycerini.....	6
Aquæ aurantii florum fortioris.....	q. s. ad 100.

To be used as a hair wash.

(*Franke.*)

3.

Hydrargyri iodidi rubri.....	1
Hydrargyri chloridi corrosivi.....	4
Alcoholis.....	50
Aquæ destillatæ.....	q. s. ad 200.

Use once daily in cases of children who cannot bear chrysarobin.

(*Quinquand.*)

4.

Naphtolis.....	1
Balsami Peruviani.....	3
Sulphuris præcipitati.....	15
Spiritus lavandulæ.....	25
Olei cadini.....	25
Linimenti saponis mollis.....	q. s. ad 100.

Apply once daily with a brush.

## 5.

Tincturæ cantharidis.....	6
Tincturæ capsici.....	6
Olei ricini.....	6
Tincturæ nucis vomicæ.....	12
Spiritus odorati.....	q. s. ad 100.

(Jamieson.)

## 6.

Cupri sulphatis.....	3
Hydrargyri ammoniati.....	3
Sulphuris sublimati.....	4
Olei cadini.....	25
Petrolati mollis.....	q. s. ad 100.

To be used once daily in cases of children who cannot bear chrysarobin.

## 7.

Acidi carbolici.....	2
Olei cadini.....	20
Sulphuris sublimati.....	20
Tincturæ iodidi.....	20
Petrolati mollis.....	q. s. ad 100.

Use as in No. 6.

(Smith.)

## 8.

Acidi borici.....	1
Acidi salicylici.....	5
Chrysarobini.....	15
Petrolati mollis.....	q. s. ad 100.

Apply once daily. Use only in severe cases occurring with older children.

(Quinquand.)

## SCABIES.

## 1.

Styracis.....	25
Balsami Peruviani.....	25
Olei ricini.....	q. s. ad 100.

Rub into the affected parts.

(Baginsky.)



2.

Creolini . . . . .	5
Olei olivæ . . . . .	q. s. ad 100.

Apply copiously over night.

(*Jacobi.*)

3.

Balsami Peruviani . . . . .	50
Glycerini . . . . .	q. s. ad 100.

Bathe the body with soft soap and warm water. Paint on the fluid and leave it on over night. Apply again next morning, omitting the bathing.

(*Monti.*)

4.

Naphtolis . . . . .	1
Cretæ præparatæ . . . . .	12
Saponis mollis . . . . .	60
Adipis . . . . .	q. s. ad 200.

Apply to the affected parts without a preceding bath. Wool-len underclothing is to be worn, or the patient should lie between woollen blankets. Powder the skin thoroughly, and do not bathe for one week.

(*Kaposi.*)

5.

Olei lavandulæ florum . . . . .	1
Olei caryophylli . . . . .	1
Potassii carbonatis . . . . .	25
Sulphuris præcipitati . . . . .	q. s. ad 100.
Adipis . . . . .	q. s. ut f. unguent.

(*Hebra.*)

6.

Sulphuris præcipitati . . . . .	10
Balsami Peruviani . . . . .	10
Petrolati mollis . . . . .	q. s. ad 100.

Bathe with soap and hot water, and apply the ointment. Repeat for two or three successive nights. The bedding and patient's clothing are to be sterilized by dry or moist heat.

(*Holt.*)

7.

Balsami Peruviani.....	50
Unguenti aquæ rosæ.....	q. s. ad 100.

Wash the skin well and rub in at night. Bathe on the following morning. Repeat daily for three or four days.

(*Widerhofer.*)

8.

Cretæ præparatæ.....	8
Olei cadini.....	15
Sulphuris sublimati.....	15
Saponis mollis.....	30
Petrolati mollis.....	q. s. ad 100.

To be used in cases in which styrax or naphthol is contra-indicated.

(*Williamson.*)

## SEBORRHŒA.

### SEBORRHŒA CAPITIS.

1.

Ætheris.....	4
Sodii boratis.....	5
Aquæ.....	q. s. ad 100.

Shake before using.

(*Hillairet.*)

2.

Glycerini.....	2
Tincturæ benzoini.....	2
Acidi borici.....	4
Alcoholis.....	q. s. ad 100.

Apply daily or every other day with a brush.

(*Kaposi.*)

3.

Sodii bicarbonatis.....	10
Aquæ.....	q. s. ad 100.

Apply after removal of the scales.

4.

Balsami Peruviani.....	2
Olei olivæ.....q. s. ad 100.	

Remove the crusts and apply locally.

5.

Acidi carbolici.....	1
Olei olivæ.....q. s. ad 200.	

Apply as in No. 4.

(*Gaucher.*)

6.

Resorcini.....	2
Unguenti aquæ rosæ.....q. s. ad 100.	

Keep the scalp covered with the ointment.

7.

Sulphuris præcipitati.....	12
Adipis lanæ hydrosi.....q. s. ad 100.	

Apply as in No. 6.

(*Holt.*)

8.

Unguenti hydrargyri oxidi flavi.....	5
Petrolati mollis.....q. s. ad 100.	

(*Ashby & Wright.*)

#### SEBORRHŒA SQUAMOSA NEONATORUM.

The body should be vigorously rubbed with oil and methodically wrapped in cloths smeared with a bland ointment. Warm baths are to be given daily and oily substances again applied. An incubator should be used, or the child should be wrapped in down or woollen coverings.

## SEBORRHŒA BALANITIS AND POSTHITIS.

1.

Amyli..... 50

Talcī.....q. s. ad 100.

Apply to the affected parts on pieces of linen or cotton.

2.

Kaolini..... 50

Magnesiæ carbonatis.....q. s. ad 100.

Apply as in No. 1.

3.

Plumbi acetatis..... 1

Aquæ.....q. s. ad 100.

Apply as in No. 1.

(Kaposi.)

---

 SYPHILIS.

## SYPHILIS HÆREDITARIA.

## INTERNAL MEDICATION.

1.

Hydrargyri chloridi mitis.....q. s.

Give from one-twelfth to one-sixth of a grain, mixed with a little sugar of milk, in a small spoonful of water, two or three times daily. Stir just before giving. If there is any tendency to diarrhœa, one-sixtieth to one-twelfth of a grain of opium may be added. Widerhofer combines from one and one-half to three grains of the saccharated carbonate of iron with the mercury.

2.

Hydrargyri tannatis.....q. s.

Give in same doses as the calomel, but omit the sugar of milk. Especially useful with children who have a tendency to diarrhœa. Opium may also be added in the same amounts as in No. 1.

(Heubner.)

## 3.

Hydrargyri iodidi flavi .....q. s.

Give from one-sixth to one-twelfth of a grain two or three times daily. Potassium iodide or any other iodide must not be given at the same time, as a caustic iodide of mercury may thus be formed in the intestines.

(Fournier.)

## 4.

Hydrargyri cum creta .....q. s.

Give from one-sixth to one-third of a grain three times a day.

(Hutchinson.)

## INUNCTIONS.

## 1.

Unguenti hydrargyri cum adipe lanæ hydroso sive  
cum unguento aquæ rosæ parati .....q. s.

Rub in from five to seven grains under light pressure for from ten to twenty minutes. This is best done in the evening. Cover the part rubbed somewhat more warm than the rest of the body, and wash it softly but thoroughly the next morning with soap and warm water. Change the place of inunction each time to a different part of the extremities and trunk. A pause of a few days is made after six inunctions.

(Heubner.)

## 2.

Unguenti hydrargyri .....q. s.

Rub the ointment into the palms of the hands and soles of the feet every morning and evening. Use five grains for each application. Otherwise apply as in No. 1.

(Hutchinson.)

## 3.

Hydrargyri oxidi rubri ..... 20  
Olei olivæ .....q. s. ad 100.

Rub in from five to fifteen grains daily. Otherwise apply as in No. 1.

(Monti.)



## 4.

Unguenti hydrargyri ..... 50

Petrolati mollis.....q. s. ad 100.

Rub in twenty grains daily. Jacobi recommends the wearing of a soft piece of leather over the knee, underneath which the ointment is applied. If preferred it may be applied under a flannel abdominal band.

## INJECTIONS.

## 1.

Hydrargyri chloridi corrosivi..... 1

Aquæ destillatæ.....q. s. ad 500.

Inject twice daily seven and one-half minims into the muscular tissues of the gluteal region. The syringe is to be directed at right angles to the skin, which should be previously antiseptically prepared. Avoid moistening with the sublimate solution the point of puncture and the opening into the tissues made by the needle.

(*Lewin.*)

## 2.

Hydrargyri chloridi corrosivi..... 1

Sodii chloridi..... 4

Aquæ destillatæ.....q. s. ad 100.

In cases of nursing children inject from one-quarter to one-half of a small hypodermatic syringe-ful. One syringe-ful may be used with larger children.

(*Monti.*)

## 3.

Hydrargyri chloridi corrosivi..... 1

Aquæ destillatæ.....q. s. ad 300.

Inject once or twice daily in doses containing from one-one-hundredth to one-fiftieth of a grain of the sublimate.

(*Jacobi.*)

## BATHS.

## 1.

Hydrargyri chloridi corrosivi..... 0.5 (gr. vijss.)

Ammonii chloridi..... 1.0 (gr. xv.)

Aquæ.....q. s. ad 100.0. (℥ iij.)

Add to one bath. As a rule one bath every other day.

(*Widerhöfer.*)

## 2.

Hydrargyri chloridi corrosivi .....	1.0 (gr. xv.)
Ammonii chloridi .....	6.0 (3 iss.)
Aquæ .....	q. s. ad 200.0. ( $\frac{2}{3}$ vjss.)

Add to one or two baths of five gallons of water.

(*Monti.*)

## 3.

Ammonii chloridi .....	2.0 (gr. xxx.)
Hydrargyri chloridi corrosivi .....	2.0. (gr. xxx.)

Add to bath of from five to seven gallons of water.

(*Comby.*)

## 4.

Hydrargyri chloridi corrosivi .....	1.0. (gr. xv.)
-------------------------------------	----------------

Add to one bath.

(*Jacobi.*)

The baths are given every day or every other day in an ordinary baby's bath tub, containing from five to seven gallons of water at a temperature of 97° F. To each bath is added from one-half a gramme to two grammes (seven and one-half to thirty grains) of corrosive sublimate, either pure or with the addition of some ammonium chloride. The bath should last from ten to fifteen minutes. It is especially indicated in cases of general eruption, although Baginsky uses this method in preference to all others for all cases. Great care is to be exercised during the bath to prevent the child from swallowing any of the contents of the tub.

## PLASTERS.

## 1.

Emplastri hydrargyri .....	q. s.
----------------------------	-------

The plaster should contain from ten to fifteen per cent of mercury. Apply a piece four inches long by two and one-half inches wide on the breast or back. Change weekly. This is the favorite method used by Fronz in Widerhofer's clinic in Vienna, with very good results.

## 2.

Olei ricini .....	7
Hydrargyri chloridi mitis vap. parat.....	23
Emplastri plumbi.....	.q. s. ad 100.

Apply a piece five inches long by two and one-half inches wide. Renew weekly.

(*Quinquand.*)

## INDIRECT AND OTHER METHODS OF TREATMENT.

It is doubtful whether the mammary glands excrete the mercury in the breast milk. This method of treating a child through the milk of the mother is too uncertain and indefinite to be of any practical value. Various other baths, injections, and preparations have been from time to time recommended, but those mentioned here may be regarded as among the most trustworthy. Injections of calomel, peptonate of mercury, and oleum cinereum are worthy of mention in addition to the sublimate, which, however, is to be preferred to all others. Fumigations with calomel and cinnabar are now very little used. The value of serotherapy and injections of syphilitic antitoxin has not yet been determined sufficiently enough to recommend their general use.

## DURATION OF TREATMENT.

No definite time can be stated. Each case differs. Hutchinson continues the treatment only while the clearly apparent symptoms are present. Hirschberg employs it through the entire first year. Holt treats a cachectic child whose parents are syphilitic, even though no active symptoms are present. He continues treatment for six weeks only, unless symptoms arise. Rotch recommends that mercury should be given a syphilitic child during the first three or four years of life at intervals of three or four months, even if no symptoms arise. He gives it at the second dentition and at puberty also. Heubner suggests that treatment should be continued for fourteen days after the last apparent symptoms have disappeared, to be repeated a year later, provided no relapse has occurred in the interim. Each relapse must be treated in the same way as the primary disease. In addition to the general treatment, local applications are of the utmost importance. These are the same as for adults. The importance of cleanliness, good air, and hygienic surroundings cannot be overestimated.

## SYPHILIS HÆREDITARIA TARDA.

## 1.

Potassii iodidi.....q. s.

Give children from the age of six upward, with affections of the bones and swelling of the liver and spleen, about fifteen grains daily. At puberty, or when the symptoms are especially severe, from twenty to thirty grains daily may be given.

(Heubner.)

## 2.

Iodoformi..... 10  
Collodii.....q. s. ad 100.

Apply locally in cases of periosteal and osteitic gummata and joint affections. Tincture of iodine, iodized glycerine, and iodol may be used in the form of solutions or ointments.

(Kaposi.)

## 3.

Potassii sive sodii iodidi..... 75  
Extracti belladonnæ foliorum alcoholici..... 1  
Aquæ destillatæ.....q. s. ad 2000.

Give from one to two teaspoonfuls three times daily after meals. The sodium salt is often better borne than the potassium.

## 4.

Unguenti hydrargyri.....q. s.

Inunctions are to be preferred to all other forms of treatment in most forms of the functional nervous disturbances, and particularly in the rapid or extensive destruction of the soft parts or bones. An inunction of twenty grains may be prescribed for children of from eight to twelve years of age, and of thirty-five grains for older children. The ointment is rubbed into both calves, or both thighs, or both arms, or the entire anterior or posterior surface of the trunk. Each cure, therefore, lasts five days and is repeated as often as is necessary. Sometimes improvement stops, when it is necessary to cease the medication for some weeks and begin anew. Iodine and mercury, either in combination or alternately, are indicated in all forms of visceral syphilis.

(Rabl.)

## GENERAL TREATMENT.

Everything should be done to strengthen the organism and overcome the cachexia, as in the early hereditary syphilis. Baths in spring water containing iodine salts, as well as the drinking of the water, are often beneficial. Iron also may be given. When the cases do not do well under the treatment outlined here, Fowler's solution in doses of from five to ten drops daily after meals may be given. The Roncegno or Levico waters may also be given in doses of from one to two teaspoonfuls noon and evening in claret with the meals. Rabl gives either of these to children who drink the iodine waters in the morning.

## ACQUIRED SYPHILIS IN CHILDREN.

Every prophylactic measure possible should be taken. Nurses and servants should be carefully examined. Drinking out of public cups, etc., should be absolutely forbidden, and above all indiscriminate kissing. The treatment is the same as that of hereditary syphilis.

---

 URTICARIA.

## 1.

*Pilocarpinæ hydrochloratis* ..... q. s.

Give in distilled water at bedtime from one-twentieth to one-eighth of a grain to a child one year old, and from one-fifteenth to one-sixth of a grain to a child from two to three years old. Gradually increase the dose.

(*Abrahams.*)

## 2.

*Atropinæ sulphatis*..... q. s.

Give one-two-hundred-and-fiftieth of a grain in the form of a powder in the evening.

(*Fräntzel.*)

## 3.

*Atropinæ sulphatis*..... 1

*Antipyrini*..... q. s. ad 100.

Give one-half of a grain of this powder in the evening.



4.

Sodii salicylatis.....q. s.

Give from five to twenty grains every two hours for three doses. Only to be used in severe cases with older children.

(Lassar.)

5.

Calcii chloridi.....q. s.

Give in doses of from fifteen to thirty grains three times daily. Get the patient under the effects of the drug rapidly, and then gradually diminish the dose.

(Wright.)

6.

Sodii chloridi.....q. s.

Moisten the affected area with cold water and rub for ten or fifteen seconds with ordinary table salt applied to the moistened end of the finger. With children, if there is much irritation, a little zinc oxide ointment may be afterward applied.

(Berliner.)

7.

Zinci oxidi..... 50

Amyli.....q. s. ad 100.

Add from one-half to one per cent of finely powdered camphor to this if the itching is very severe.

(Gaucher.)

8.

Bismuthi salicylatis..... 10

Amyli.....q. s. ad 100.

9.

Dermatolis ..... 10

Magnesiæ.....q. s. ad 100.

10.

Hydrargyri chloridi corrosivi..... 1

Acidi hydrochlorici..... 1

Aquæ laurocerasi .....100

Aquæ destillatæ .....q. s. ad 2500.

(Brocq.)

## 11.

Chloralis .....	1
Glycerini .....	5
Aquæ.....	q. s. ad 100.

## 12.

Glycerini .....	40
Balsami Peruviani.....	q. s. ad 100.

*(Monti.)*

## 13.

Glycerini .....	3
Ætheris .....	5
Alcoholis.....	q. s. ad 200.

## 14.

Tincturæ benzoini.....	3
Mentholis .....	5
Glycerini.....	5
Alcoholis.....	q. s. ad 200.

*(Kaposi.)*

## 15.

Ichthyolis.....	1
Petrolati mollis .....	q. s. ad 100.

## 16.

Unguenti hydrargyri ammoniati.....	20
Petrolati mollis .....	q. s. ad 100.

*(Holt.)*

## 17.

Mentholis .....	1
Zinci oxidi.....	10
Petrolati mollis.....	q. s. ad 100.

## 18

Acidi carbolicæ.....	1
Bismuthi subnitratæ.....	10
Petrolati mollis.....	q. s. ad 100.

*(Gaucher.)*

## VERRUCÆ.

## 1.

Acidi nitrici..... q. s.

Apply alone on a small piece of wood, or immediately follow it by the application of a few drops of pure liquid carbolic acid. The latter method is recommended by Laubenberg.

## 2.

Hydrargyri chloridi corrosivi..... 15  
 Alumini..... 15  
 Plumbi carbonatis..... 15  
 Camphoræ ..... 15  
 Acidi acetici diluti ..... 15  
 Alcoholis.....q. s. ad 100.

Paint the surface.

(*Plenck*)

## 3.

Glycerini..... 20  
 Acidi acetici glacialis..... 30  
 Sulphuris sublimati .....q. s. ad 100.

Apply daily to each wart. Continue treatment for several days. Shake well before using. To be used in cases of multiple warts of the face in older children.

## 4.

Acidi acetici glacialis..... 8  
 Sulphuris sublimati..... 15  
 Glycerini .....q. s. ad 100.

To be used in cases in which No. 3 is too strong.

(*Kaposi.*)

## 5.

Acidi salicylici..... 5  
 Collodii ..... 15  
 Ætheris.....q. s. ad 100.

Paint the surface.

(*Shoemaker.*)

6.

Acidi salicylici ..... 12

Collodii.....q. s. ad 100.

Apply twice a day for three days, after which soak the part in warm water, and use pumice soap if not inflamed. Repeat if necessary.

(Rotch.)

## ZOSTER

1.

Spiritus camphoræ.....q. s.

Give five drops on a lump of sugar three times daily.

(Drinkwater.)

2.

Liquoris potassii arsenitis .....q. s.

Give in gradually increasing doses for the neuralgia continuing after the disappearance of the eruption. If the pain is very severe, either morphine, opium, or chloral may be used in older children.

(Kaposi.)

3.

Salolis..... 25

Ætheris.....q. s. ad 100.

Apply with a brush twice daily.

(Bilkstein.)

4.

Opii pulveris ..... 1

Camphoræ ..... 2

Zinci oxidi..... 20

Amyli..... q. s. ad 100.

(Robin.)

5.

Extracti opii.....	1
Cerati .....	100.

Not to be used with very young children.

6.

Cocainæ hydrochloratis.....	1
Unguenti .....	q. s. ad 100.

Not to be used over a large area with very young children.

(*Kaposi.*)

7.

Acidi carbolici.....	1
Mentholi .....	1
Unguenti zinci oxidi.....	50
Unguenti hydrargyri ammoniati.....	q. s. ad 100.

Mix and make stiff by adding zinc oxide. Apply thickly on a jacket of lint.

(*Taylor.*)





# INDEX.

## A.

- Acarus—see scabies, 92
- Achorion, 17
- Acne, treatment, 95
  - disseminata, 95
  - vulgaris, 95
  - treatment—see acne, 95
- Acné varioliformis—see molluscum contagiosum, 87
- Acquired syphilis in children, treatment, 154
- Alligator boy, 38
- Alopecia areata, 1
  - diagnosis, 2
  - etiology, 6
  - prognosis, 9
  - symptoms, 2
  - treatment, 6, 97
- Alopecia areata, plate, 1
- Alopecia circumscripta—see alopecia areata, 1
- Antiseptic cage treatment of burns, 100
- Antistreptococcic serum treatment of erysipelas, 118
- Area Celsi—see alopecia areata, 1

## B.

- Bacillus tuberculosis, 68, 72
- Baldness in contagious impetigo, 24
- Baths in eczema, 49
  - in syphilis, 49
- Birthmark, 61
- Bites of insects, treatment, 99
- Boil, treatment—see furuncle, 123
- Burns, treatment, 100
  - powder, treatment, 101

## C.

- Chilblains, treatment—104
- Chloasma, treatment—see freckles and other pigmentations, 122
- Chromophytosis, 11
- Claret stain—see nævus flammeus, 61
- Clothes louse, treatment—see phthiriasis corporis, 138
- Combustio, treatment—see burns, 100
- Condyloma subcutaneum—see molluscum contagiosum, 87
- Congenital ichthyosis, 37
- Contagious impetigo, 21, 46
  - diagnosis, 21
  - etiology, 21
  - prognosis, 25
  - symptoms, 21
  - treatment, 25, 128
  - on the scalp, 24
- Crab louse, treatment—see phthiriasis pubis, 138
- Crusta lactea, 21
- Crusted ringworm—see favus, 121

## D.

- Dental burr in treatment of lupus vulgaris, 72
- Dentition, influence of, in etiology of eczema, 47
- Dermatitis contusiformis—see erythema nodosum, 84, 120
- exfoliativa, 28
- exfoliativa neonatorum, treatment, 106
- gangrænosa, treatment, 105

Dermatitis gangrænosa infantum,  
treatment—see dermatitis gan-  
grænosa, 105

Diffuse congenital keratoma, 37

Driving in of eczema, 47

## E.

Ecthyma, 23

Ectropion in lupus, 70

Eczema, 41

etiology, 47

symptoms, 47

treatment, 48

acute, 41

treatment, 48, 106

ani, treatment, 113

aurium, treatment, 113

capitis, treatment, 114

chronic, 42

treatment, 48, 110

erythematosum, 43

impetiginosum, 46

intertrigo—see erythema inter-  
trigo, 43

treatment, 129

labiorum, treatment, 116

narium, treatment, 116

of the nails, treatment, 134

palpebrarum, treatment, 117

papular, 44

pustular, 46

rubrum, 46

squamous, 46

umbilici, treatment, 117

Eczema, plate, 41

Ephelides—see freckles, 122

Erysipelas, treatment, 118

Erythema centrifuge—see lupus  
erythematosus, 72

Erythema, treatment, 119

intertrigo, 43

multiforme—see erythema no-  
dosum, 84

nodosum, 84

diagnosis, 84

etiology, 84

prognosis, 85

symptoms, 84

treatment, 85, 120

pernio, treatment—see chil-  
blains, 104

Erythematous eczema, 43

syphilide, 83

Excessive sweating, treatment—  
see hyperidrosis, 124

## F.

Favus, 11

of nails, treatment, 135

on hairy parts, 15

diagnosis, 15

etiology, 17

prognosis, 19

prophylaxis, 19

symptoms, 15

treatment, 18, 121

on non-hairy parts, 12

diagnosis, 12

etiology, 17

prognosis, 18

prophylaxis, 19

symptoms, 12

treatment, 18, 121

Fish-skin disease, 35

Folliculitis decalvans, 1

Formulary and note, 95

Freckles, treatment, 122

Furuncle, treatment, 123

Furunculus, treatment—see furun-  
cle, 123

## G.

Galvano-cauterization treatment of  
furuncle, 124

Guttate form of psoriasis, 27

## H.

Harlequin foetus, 37

Head louse, treatment—see phthiri-  
asis capitis, 137

Hemorrhage, idiopathic, 85

Hereditary syphilis, 83

Herpes esthiomenos—see lupus, 67

tonsurans—see ringworm upon  
hairy parts, 13

zoster, treatment—see zoster, 158

Hives, treatment—see zoster, 158

Honeycomb ringworm—see favus, 11

Hyperidrosis, treatment, 124

## I.

- Ichthyose—see ichthyosis, 35  
 Ichthyosis, 35  
     diagnosis, 35  
     etiology, 40  
     prognosis, 35  
     symptoms, 35  
     treatment, 40  
     congenita, 37  
     hystrix, 37, 51  
     linearis neuropathica localis, 51  
     mitis, 35  
     simplex, 35  
     vera—see ichthyosis, 35  
 Ichthyosis, plate, 35  
 Idiopathic hemorrhage, 85  
 Idrosis, treatment—see hyperidrosis, 124  
 Ignis sacer treatment—see zoster, 158  
 Impetigo—see contagious impetigo, 21  
     treatment, 25, 128  
     contagiosa—see contagious impetigo, 21  
 Impetigo contagiosa, plate, 21  
 Infantile syphilis, hereditary, 83  
 Intertrigo, treatment, 129  
 Itch, treatment—see scabies, 92

## K.

- Kelis—see keloid, 90  
 Keloid, 90  
     diagnosis, 91  
     etiology, 90  
     prognosis, 92  
     symptoms, 90  
     treatment, 92  
     false, 90  
     true, 90  
 Keratosis follicularis, 89  
     diagnosis, 89  
     etiology, 89  
     prognosis, 90  
     symptoms, 89  
     treatment, 90  
     follicularis, Darier, 90  
     pilaris, 89

## L.

- Lentigo, treatment—see freckles, 122  
 Lice, treatment—see phthiriasis, 137

- Lichen pilaris, 89  
     planus, 80, 83  
         diagnosis, 80  
         prognosis, 82  
         symptoms, 80  
         treatment, 82  
 Lichen ruber, 75  
     diagnosis, 75  
     prognosis, 79  
     symptoms, 75  
     treatment, 80

## forms of:

- papular, 78  
 rugous, 79  
 squamous, 78  
 Lichen ruber, plate, 75  
 Lichen simplex—see papular eczema, 44  
     tropicus, treatment—see miliaria, 133  
 Linear papilloma—see papilloma lineare, 61  
 London hospital treatment of burns, 100  
 Lupus, 67

- diagnosis, 68  
     etiology, 67  
     symptoms, 68  
     treatment, 72, 130  
     and other tuberculides, 67  
     erythematodes—see lupus erythematosus, 72  
     erythematosus, 72  
         treatment, 72, 132  
     superficialis—see lupus erythematosus, 72  
     verrucosus—see tuberculosis verrucosa, 70  
     vulgaris—see lupus, 67  
 Lupus vulgaris, plate, 67

## M.

- Miliaria, treatment, 133  
     alba,  
     rubra,  
         treatment—see miliaria, 133  
 Molluscum, 87  
     contagiosum, 87  
     diagnosis, 89  
     etiology, 88

- Molluscum contagiosum, prognosis, 89  
 symptoms, 87  
 treatment, 89  
 fibrosum, 87  
 sebaceum, 87  
 Morbus pedicularis, treatment—see  
 phthisiasis, 137
- N.
- Nævi, pigmented and hairy, 55  
 Nævus araneus, 61  
   treatment, 64, 134  
 cavernosus, 61, 63  
   treatment, 64, 134  
 flammeus, 61  
   treatment, 64, 134  
 pigmentosus, 55  
   treatment, 57  
 pilosus, 55  
   treatment, 58  
 sanguineus—see vascular nævus, 61  
 tuberosus, 61, 62  
   treatment, 65, 134  
 unius lateris, 51, 52  
 vascularis—see vascular nævus, 61  
 verrucosus, 37, 51  
 Nævus pilosus, plate, 55  
 Nails, diseases of, treatment :  
   eczema, 134  
   favus, 135  
   psoriasis, 135  
   ringworm, 135, 136  
 Nerve nævus—see papilloma lineare, 51  
 Nettle rash, treatment—see urticaria, 154
- O.
- Onychomycosis favosa, treatment—  
 see favus of the nails, 135  
 trichophytina, treatment—see  
 ringworm of the nails, 136
- P.
- Papilloma lineare, 51  
   diagnosis, 51  
   symptoms, 51  
   treatment, 53  
 Papilloma lineare, plate, 51  
   neuroticum—see papilloma lineare, 51  
 Papular eczema, 44  
   syphilide, 30, 83  
 Parchment skin, 35  
 Paronychia, in contagious impetigo, 25  
 Pediculosis, treatment:  
   capitis, 137  
   corporis, 137  
   pubis, 136  
     see phthiriasis, 137  
 Pediculus vestimentorum, treatment—see phthiriasis corporis, 138  
 Peliosis rheumatica, 87  
 Pemphigus gangrænosus, treatment—see dermatitis gangræno-  
 sa, 105  
 Pernio, treatment—see chilblains, 104  
 Phtheiriasis, treatment—see phthiriasis, 137  
 Phthiriasis, treatment:  
   capitis, 137  
   corporis, 138  
   pubis, 138  
 Pigmentary nævus, 55  
 Pityriasis, 43  
   pilaris—see keratosis follicularis, 89  
   rubra pilaris, 75  
 Porcupine skin, 37  
 Porrigo contagiosa—see contagious  
 impetigo, 21  
   decalvans—see alopecia areata, 1  
   furfurans—see ringworm upon  
   hairy parts, 13  
 Portwine mark, 61  
 Powder burns, treatment, 101  
 Prickly heat, treatment—see mili-  
 aria, 133  
 Prurigo, treatment, 139  
 Pruritus cutaneus, treatment, 140  
 Psoriasis, 27  
   diagnosis, 27  
   etiology, 32  
   prognosis, 32  
   symptoms, 27



- Psoriasis, treatment, 32, 141  
 forms of :  
   guttate, 27  
   nummular, 27  
   punctate, 27  
   of the nails, treatment, 135  
 Psoriasis, plate, 27  
 Purpura, 85  
   diagnosis, 87  
   etiology, 85  
   prognosis, 86  
   symptoms, 85  
   treatment, 87  
   hæmorrhagica, 86  
   papulosa, 86  
   rheumatica, 87  
     treatment, 142  
 Pus cocci, in contagious impetigo, 21  
 Pustular eczema, 21
- Q.
- Quinine, in erythema nodosum, 120
- R.
- Ringworm and favus, 11  
 Ringworm of the nails, treatment, 136  
   of the scalp—see ringworm  
   upon hairy parts, 13  
   upon hairy parts, 13  
     diagnosis, 13  
     etiology, 17  
     prognosis, 19  
     prophylaxis, 19  
     symptoms, 13  
     treatment, 17, 143  
   upon non-hairy parts, 12  
     diagnosis, 12  
     etiology, 17  
     prognosis, 18  
     prophylaxis, 19  
     symptoms, 12  
     treatment, 18
- S.
- Scabies, 92  
   diagnosis, 92  
   etiology, 92  
   prognosis, 93  
   symptoms, 92  
   treatment, 93, 144
- Scrofulide érythémateuse—see lupus erythematosus, 72  
 tuberculeuse—see lupus, 67  
 Scrofuloderma, 71  
 Seborrhœa, balanitis and posthitis, treatment, 148  
   capitis, treatment, 146  
   squamosa neonatorum, treatment, 147  
 Shingles, treatment—see zoster, 158  
 Stings of insects, treatment, 99  
 Strumous dactylitis, 71  
 Sudamina, treatment—see miliaria, 133  
 Sweating, excessive, treatment—see hyperidrosis, 124  
 Syphilides, forms of:  
   bullous, 83  
   erythematous, 83  
   papular, 83  
   pustular, 83  
 Syphilis, acquired, in children, treatment, 154  
   hæreditaria, 83  
     diagnosis, 83  
     symptoms, 83  
     treatment, 84, 148  
       (a) baths, 150  
       (b) duration of, 152  
       (c) general, 84  
       (d) indirect and other methods of, 152  
       (e) injections, 150  
       (f) internal medication, 84, 148  
       (g) inunctions, 84, 149  
       (h) local, 84, 152  
       (i) plasters, 151  
       (j) prophylaxis, 83  
 Syphilis hæreditaria, plate, 83  
 Syphilis hæreditaria tarda, treatment, 153
- T.
- Telangiectasis, 61  
 Tetter—see eczema, 41  
 Tinea circinata—see ringworm upon non-hairy parts, 12  
   decalvans—see alopecia areata, 1  
   favosa—see favus, 11

*Tinea favosa unguium*, treatment—  
 see favus of the nails, 135  
*furfuracea*, treatment—see se-  
 borrhoea, 146  
*kerion*—see kerion, 14  
*lupinosa*—see favus, 11  
*tonsurans*—see ringworm upon  
 hairy parts, 13  
*trichophytina*—see ringworm,  
 11  
*trichophytina unguium*, treat-  
 ment—see ringworm of the  
 nails, 136  
*versicolor*, 11  
 Treatment of powder burns, 101  
*Trichophyton*, 17  
*Trichophytosis*, 11  
*Trichophytosis capitis*, plate, 11  
*Tuberculides*, lupus and other, 67  
*Tuberculosis bacillus*, in lupus, 68  
 in lupus erythematosus, 72  
*Tuberculosis verrucosa*, 70

## U.

Ulcers, of conjunctiva and buccal  
 mucous membrane, in impetigo,  
 25  
*Urticaria*, treatment, 154

## V.

Vaccination, double infection with,  
 24  
 Vaccine-lymph treatment of *nævus*  
*vasculosus*, 134  
*Varicella gangrænosa*, treatment—  
 see dermatitis gangrænosa, 105  
 Various skin diseases of children,  
 83  
 Vascular *nævus*, 61  
 Vascular *nævus*, plate, 61  
 Vegetable parasites, 11  
*Verruca*, 51  
 treatment, 157

## W.


Warts, 51  
 treatment—see *verrucae*, 157  
 Water-bed treatment of burns, 101

## X.

Xeroderma, 35  
*ichthyoides*—see *ichthyosis*, 35

## Z.

*Zona*, treatment—see *zoster*, 158  
*Zoster*, treatment, 158





# MEDICAL PUBLICATIONS

OF

## WILLIAM WOOD & COMPANY.

---

**Abney, Capt. W. de W., C.B., D.C.L., F.R.S.,**

Late Royal Engineers.

COLOUR VISION, being the Tyndall Lectures delivered in 1894 at the Royal Institution. One volume, 8vo, 241 pages, illustrated by a chromo-lithographic plate and numerous diagrams, muslin, \$2.00.

**Adams, Francis, LL.D., Surgeon.**

THE GENUINE WORKS OF HIPPOCRATES. Translated from the Greek, with a Preliminary Discourse and Annotations. 8vo, 766 pages, gilt top, extra muslin, \$5.00.

**Allingham, William, F.R.C.S. Lond.**

and

**Allingham, Herbert W., F.R.C.S. Lond.**

THE DIAGNOSIS AND TREATMENT OF DISEASES OF THE RECTUM. Being a Practical Treatise on Fistula, Piles, Fissure and Painful Ulcer, Procidencia, Polypus, Stricture, Cancer, etc. Sixth Edition. One volume of 500 pages, 8vo, with numerous illustrations. Cloth, \$4.00.

**American Journal of Obstetrics and Diseases of Women and Children.**

Issued monthly. Price, \$5.00 a year.

**Anatomical Remembrancer ;**

OR, COMPLETE POCKET ANATOMIST; containing a concise description of the bones, ligaments, muscles, and viscera; the distribution of the nerves, blood-vessels, and absorbents; the arrangement of the several fasciæ; and the organs of the senses. (Wood's Pocket Manuals.) 18mo, 297 pages, muslin, \$1.00.

**Ball, James B., M.D. Lond., M.R.C.P.**

A HAND-BOOK OF DISEASES OF THE NOSE AND PHARYNX. Second revised edition. One volume of 376 pages, 12mo, illustrated, muslin, \$2.50.



**Baruch, Simon, M.D.,**

New York City.

ON HYDROTHERAPY. One volume, 8vo. (In preparation.)

**Bell, A. N., A.M., M.D.,**

Editor of "The Sanitarian;" Member of American Medical Association, American Public Health Association, Medical Society of the State of New York; Honorary Member of Connecticut Medical Society; Corresponding Member of the Epidemiological Society of London; formerly P. A. Surgeon, U. S. Navy, etc.

CLIMATOLOGY OF THE UNITED STATES AND ADJACENT COUNTRIES, and of such Foreign Ports and Places as have Intimate Commercial Relations with the United States, with special reference to Health Resorts, and the Protection of Public Health. 8vo, 393 pages, muslin, \$1.00.

**Benedikt, Moriz,**

Professor at Vienna.

ANATOMICAL STUDIES UPON BRAINS OF CRIMINALS. A CONTRIBUTION TO ANTHROPOLOGY, MEDICINE, JURISPRUDENCE, AND PSYCHOLOGY. Translated from the German by E. P. FLOWER, M.D., New York; Department of Translation, New York Medico-Chirurgical Society. Illustrated with wood-engravings. One volume, 8vo, 185 pages, muslin, \$1.50.

**Blyth, A. W., M.R.C.S.**

POISONS, THEIR EFFECTS AND DETECTION. A manual for the use of analytical chemists and experts. With an introductory essay on the Growth of Modern Toxicology. Volume I., with tables and illustrations. Volume II., with tables and illustrations. 2 vols., 8vo, 699 pages, muslin, \$2.00.

**Bock, C. E., M.D.,**

Professor at Leipsic.

ATLAS OF HUMAN ANATOMY. With Explanatory Text. Containing thirty-eight steel plates, most of them hand-colored, of the bones, muscles, vessels, and nerves of the human body; organs of sense, eye, ear, nose, and tongue; respiratory apparatus, abdominal and pelvic viscera, organization of fœtus, the teeth, with the genito-urinary organs of the male and female. One volume, large 4to, muslin. \$15.00.

**Bodenhamer, William, M.D.,**

Professor of the Diseases, Injuries, and Malformations of the Rectum, Anus, and Genito-Urinary Organs.

TREATISE ON THE HEMORRHOÏDAL DISEASE. Giving its History, Nature, Cause, Pathology, Diagnosis, and Treatment. One volume, 8vo, over 300 pages, illustrated by two chromo-lithographic plates and many wood-cuts. Muslin, \$3.00.

**Bodenhamer, William, M.D.**

PRACTICAL OBSERVATIONS ON THE ÆTIOLOGY, PATHOLOGY, DIAGNOSIS, AND TREATMENT OF ANAL FISTURE. Illustrated by numerous cases and drawings. One volume, 8vo, 199 pages, muslin, \$2.25.

AN ESSAY ON RECTAL MEDICATION. One volume, 8vo, 58 pages, illustrated, muslin, \$1.00.

**Bosworth, Francke Huntington, M.D.,**

Professor of Diseases of Throat in Bellevue Hospital Medical College, New York; Consulting Laryngologist to the Presbyterian and St. Vincent Hospitals, New York; Honorary Fellow of the British Otological, Laryngological, and Rhinological Association; Corresponding Member of the Société Française d'Otologie et de Laryngologie; Fellow of the American Laryngological Association, the Academy of Medicine, New York; Member of the Medical Society of New York State, and of the County Society of New York, etc.

A TEXT-BOOK ON DISEASES OF THE NOSE AND THROAT. Complete in one volume, 834 pages, octavo, numerous wood-engravings, and full-page chromo-lithographic plates. Muslin, \$5.50; leather, \$6.50.

DISEASES OF THE THROAT AND NOSE. In two volumes, 8vo.

Vol. I.—DISEASES OF THE NOSE AND NASO-PHARYNX. 8vo, 670 pages, 4 colored plates and 182 wood-cuts. Extra muslin, \$6.00.

Vol. II.—DISEASES OF THE THROAT. 8vo, 832 pages, 3 colored plates, and 125 wood-cuts. Extra muslin, \$6.00.

**Bradford, Edward H., M.D.,**

Surgeon to the Children's Hospital, Boston City Hospital, and Samaritan Hospital; Instructor in Clinical Surgery, Harvard Medical School; and

**Lovett, Robert W., M.D.,**

Surgeon to the Samaritan Hospital; Assistant Out-Patient Surgeon to the Children's Hospital; Out-Patient Surgeon to the Carney Hospital; formerly Assistant Surgeon to the New York Orthopedic Dispensary and Hospital.

A TREATISE ON ORTHOPEDIC SURGERY. Illustrated with seven hundred and eighty-nine wood-engravings. 8vo, 790 pages, extra muslin, \$6.00.

**Bramwell, Byrom, M.D., F.R.C.P. Ed., F.R.S. Ed.**

DISEASES OF THE SPINAL CORD. Third Edition. Illustrated by 170 engravings in black and numerous colors. In this edition the subject matter has been thoroughly revised and greatly extended; in fact, the text has been entirely rewritten and rearranged in lecture form. A large number of new illustrations—chiefly clinical—have been added. Extra muslin, octavo, 671 pages, \$5.00.

**Brass, Dr. Arnold,**

Göttingen.

ATLAS OF HUMAN HISTOLOGY. Sixty full-page plates, engraved and superbly printed in many colors, with explanatory notes. An exquisite work. Authorized translation from the German, with additions. By R. A. YOUNG, M.D., B.Sc. Lond. Extra buckram, in box, \$10.00 net.

**Broadbent, John F. H., M.D., M.R.C.P.**

ADHERENT PERICARDIUM. One 8vo volume of 126 pages, with numerous references, etc., muslin, \$1.25.

**Brouardel, P., M.D. Paris.**

DEATH AND SUDDEN DEATH. Translated by F. LUCAS BENHAM, M.D., B.S. Lond. One volume, 8vo, 280 pages, muslin, \$3.00.

**Buck, Albert H., M.D.,**

Clinical Professor of the Diseases of the Ear, in the College of Physicians and Surgeons, New York; Consulting Aural Surgeon, New York Eye and Ear Infirmary.

A MANUAL OF DISEASES OF THE EAR, for the use of Students and Practitioners of Medicine. (American Series of Medical Text-Books.) Second Edition. 468 pages, 117 illustrations, extra muslin, \$2.50.

A VEST-POCKET MEDICAL DICTIONARY. Embracing those terms and abbreviations which are commonly found in the medical literature of the day, but excluding names of drugs and many words which may more properly be found in a general dictionary of the English language. A most complete little book of 536 pages, less than one-half inch in thickness, 32mo, bound in flexible leather. Price, \$1.00.

A TREATISE ON HYGIENE AND PUBLIC HEALTH. By various authors. Edited by ALBERT H. BUCK, M.D. In two volumes, royal 8vo, 702 and 657 pages. Illustrated by numerous wood-engravings. Price, per volume, in leather, \$6.00; and in morocco, \$7.50.

**Cabot, Richard C., M.D.,**

Boston, Mass.

A GUIDE TO THE CLINICAL EXAMINATION OF THE BLOOD FOR DIAGNOSTIC PURPOSES. Second edition. One volume of 430 pages, octavo, illustrated by numerous wood-engravings and by chromo-lithographic plates, muslin, \$3.75.

**Carpenter, Wm. B., C.B., M.D., LL.D.**

THE MICROSCOPE AND ITS REVELATIONS. 8vo. Vol. I., 388 pages; Vol. II., 354 pages. One colored, twenty-six plain plates, and five hundred and two wood-engravings. Two volumes in one. Muslin, \$4.00.

**Cheyne, W. Watson, M.B., F.R.S., F.R.C.S. Lond.**

THE OBJECTS AND LIMITS OF OPERATIONS FOR CANCER, with special reference to Cancer of the Breast, Mouth, and Throat, and Intestinal Tract. Being the Lettsomian Lectures for 1896. 8vo, 146 pages, muslin, \$1.75.

**Corson, John W., M.D.,**

Late Physician to the class of "Diseases of the Chest and Throat" in the New York and Eastern Dispensaries; formerly Physician to the Brooklyn City Hospital; Physician to the Orange Memorial Hospital, etc.

ON THE TREATMENT OF PLEURISY. With an Appendix of Cases, showing the Value of Combinations of Croton Oil, Ether, and Iodine, as Counter irritants in other Diseases. 16mo, 31 pages, muslin, 50c.

**Curtis, Edward, A.M., M.D.**

MANUAL OF GENERAL MEDICINAL TECHNOLOGY, INCLUDING PRESCRIPTION WRITING. Third edition, revised in accordance with the U. S. P. for 1890. One volume, 32mo, 243 pages, muslin, \$1.00. (Wood's Pocket Manuals.)

**Dana, Charles L., A.M., M.D.,**

Professor of Nervous and Mental Diseases in the New York Post-Graduate Medical School, and in Dartmouth Medical College; Visiting Physician to Bellevue Hospital, etc.

TEXT-BOOK OF NERVOUS DISEASES. Being a Compendium for the Use of Students and Practitioners of Medicine. Fourth edition, revised and enlarged. 8vo, 640 pages, 210 illustrations, \$3.50 *net*.

**Delafield, Francis, M.D.,**

Professor of Pathology and Practical Medicine, College of Physicians and Surgeons, New York.

STUDIES IN PATHOLOGICAL ANATOMY. Volume I., treating of the following subjects: Phthisis, Peritonitis, Pleurisy, Pneumonia, Empyema, Hydrothorax, Bronchitis, and Tuberculosis. Illustrated with ninety-three full-page and double-page plates made by the following processes: Wood-engravings of Original Drawings on the Block, Etchings on Copper, Lithographs from Original Drawings on the Stone, and Photographs of Specimens. Royal 8vo, bound in half morocco, gilt top, plates hinged on linen guards, \$20.00.

Volume II.: Broncho-Pneumonia, Chronic Phthisis, Lobar Pneumonia, Acute Bright's Disease, Chronic Bright's Disease. Illustrated with one hundred and thirty-three full and double-page plates hinged on linen guards, similar to those of Vol. I. Royal 8vo, bound in half morocco, gilt top, \$20.00.

**Donkin, H. Bryan, M.D. Oxon., F.R.C.P.,**

Physician to the Westminster Hospital and to the East London Hospital for Children at Shadwell; Joint Lecturer on Medicine and Clinical Medicine at Westminster Hospital Medical School.

THE DISEASES OF CHILDHOOD (MEDICAL). 8vo, 450 pages, muslin, \$4.00.



**Delafield and****Prudden, T. Mitchell, M.D.,**

Professor of Pathology and Director of the Laboratories of Histology, Pathology, and Bacteriology, College of Physicians and Surgeons, Columbia College, New York.

**A HANDBOOK OF PATHOLOGICAL ANATOMY AND HISTOLOGY.** With an Introductory Section on Post-Mortem Examinations and the Methods of Preserving and Examining Diseased Tissues. Fifth Revised and Enlarged Edition. One volume of 846 pages, 8vo, illustrated by engravings in black and many colors, and a chromo-lithographic plate. Muslin, \$6.00; leather, \$7.00.

**Draper, John C., M.D., LL.D.,**

Professor of Chemistry in the Medical Department, University of New York, and of Physiology and Natural History in the College of the City of New York.

**A PRACTICAL LABORATORY COURSE IN MEDICAL CHEMISTRY.** One volume of 80 pages, printed on one side only, oblong, for laboratory use, muslin, \$1.00.

**Dwight, Thomas, A.M., M.D.,**

Instructor in Topographical Anatomy and Histology in Harvard University; Fellow of the American Academy of Arts and Sciences; Surgeon at Carney Hospital.

**FROZEN SECTIONS OF A CHILD.** Fifteen full-page lithographic plates, drawings from nature by H. P. QUINCY, M.D. One volume, royal 8vo, 66 pages, muslin, \$3.00.

**Einhorn, Max, M.D.,**

Instructor in Clinical Medicine, New York Post-Graduate Medical School; Visiting Physician to the German Dispensary, and to the Montefiore Home for Chronic Invalids.

**DISEASES OF THE STOMACH.** Complete in one volume of 461 pages, post-octavo (uniform with other volumes of the Medical Practitioners' Series). Muslin, \$3.50; flexible morocco, \$4.25.

**Ellis, George Viner, M.D.,**

Professor of Anatomy in University College, London; and

**Ford, G. H., Esq.**

**ILLUSTRATIONS OF DISSECTIONS.** In a series of original colored plates, representing the dissections of the human body, with descriptive letterpress. The drawings are from nature by Mr. Ford, from directions by Prof. Ellis. Containing fifty-six full-page chromo-lithographic plates. Two volumes in one, 8vo, 459 pages, muslin, \$4.00.

**Ewart, William, M.D. Cantab., F.R.C.P. Lond., M.R.C.S. Eng.,**

Physician to St. George's Hospital, and to the Belgrave Hospital for Children; formerly Assistant Physician and Pathologist to the Brompton Hospital for Consumption, etc., etc.

**THE PULSE SENSATIONS: A STUDY IN TACTILE SPHYGMOLOGY.** 8vo, 510 pages, profusely illustrated, muslin, \$4.00.

**GOUT AND GOUTINESS AND THEIR TREATMENT.** One volume of 601 pages, 8vo, muslin, \$5.00.



**Finger, Ernest, M.D.,**

Docent at the University of Vienna.

GONORRHŒA: being the translation of "Blennorrhœa of the Sexual Organs and its Complications." With seven full-page plates in colors and thirty-six wood engravings in the text. Third revised edition, 8vo, 330 pages, muslin, \$3.00.

**Fowler, E. P., M.D.**

SUPPRESSION OF URINE. Clinical Descriptions and Analysis of Symptoms. 86 pages, illustrated, muslin, \$1.50.

**Fox, George Henry, A.M., M.D.,**

Professor of Diseases of the Skin in the College of Physicians and Surgeons, Columbia University, New York, etc., etc.

SKIN DISEASES OF CHILDREN. The work is based upon a series of papers originally contributed to the *American Journal of Obstetrics*, in 1896, and has been elaborated and a large formulary added. 8vo, profusely illustrated by photogravure plates, chromo-lithographic plates, and half-tone cuts. Muslin, \$2.50 *net*.

**Frey, Heinrich,**

Professor of Medicine in the University of Zurich; Surgeon New York Eye and Ear Infirmary; Ophthalmic and Aural Surgeon to the St. Catherine and Williamsburg Hospitals, etc., etc.

THE MICROSCOPE AND MICROSCOPICAL TECHNOLOGY. A Text-book for Physicians and Students. Translated and edited by GEO. R. CUTTER, M.D. Illustrated by three hundred and eighty-eight engravings on wood. 8vo, 660 pages, muslin, \$6.00.

**Freyer, P. J., M.A., M.D., M.Ch.,**

Surgeon Lieutenant-Colonel, Bengal Army (retired).

THE MODERN TREATMENT OF STONE IN THE BLADDER BY LITHOLAPAXY. A description of the operation and instruments, with cases illustrative of the difficulties and complications met with. Second Edition. One 8vo volume, illustrated. Muslin, \$1.50.

**Garrigues, Henry Jacques, A.M., M.D.,**

Obstetric Surgeon to the Maternity Hospital; Physician to the Gynecological Department of the German Dispensary; Fellow of the American Gynecological Society; Fellow of the New York Obstetrical Society, etc.

DIAGNOSIS OF OVARIAN CYSTS BY MEANS OF THE EXAMINATION OF THEIR CONTENTS. 8vo, 112 pages, illustrated, muslin, \$1.25.

**Gemmell, G. H., F.I.C., F.C.S. Ed.**

CHEMICAL NOTES AND EQUATIONS, INORGANIC AND ORGANIC. One volume of 254 pages, 12mo. Muslin, \$1.75.

**Gowers, W. R., M.D.,**

Assistant Professor of Clinical Medicine in University College; Senior Assistant Physician to University College Hospital; Physician to the National Hospital for the Paralyzed and Epileptic.

EPILEPSY AND OTHER CHRONIC CONVULSIVE DISEASES. Their Causes, Symptoms, and Treatment. 8vo, 366 pages, muslin, \$1.00.

DIAGNOSIS OF THE DISEASES OF THE BRAIN AND SPINAL CORD. 8vo, 301 pages, muslin, \$1.00.

**Grandin, Egbert H., M.D.,**

Chairman Section on Obstetrics and Gynecology, New York Academy of Medicine; Obstetric Surgeon, New York Maternity Hospital; Obstetrician, New York Infant Asylum, etc.; and

**Gunning, Josephus H., M.D.,**

Instructor in Electro-Therapeutics, New York Post-Graduate Medical School and Hospital; Gynecologist to Riverview Rest for Women; Electro-Gynecologist, Northeastern Dispensary, etc.

PRACTICAL TREATISE ON ELECTRICITY IN GYNECOLOGY. Illustrated. 8vo, muslin, 180 pages, \$2.00.

**Haab, Prof. O., M.D.,**

Professor of Ophthalmology, University of Zurich. Translated and Edited by

**Ernest Clarke, M.D., B.S. Lond.,**

Fellow of the Royal College of Surgeons; Surgeon to the Central London Ophthalmic Hospital; Ophthalmic Surgeon to the Miller Hospital, etc., etc.

AN ATLAS ON OPHTHALMOSCOPY. One 12mo volume, containing 64 full-page plates, embracing 102 figures, superbly executed by chromo-lithography, with complete descriptive text and an introductory chapter. The plates are probably the finest set of ophthalmoscopic figures ever published. Muslin, \$3.00 *net*. (Volume I. of Wood's Medical Hand Atlases.)

**Hamilton, Frank Hastings, A.M., M.D., LL.D.,**

Professor of the Practice of Surgery, with Operations, and of Clinical Surgery, in Bellevue Hospital Medical College; Visiting Surgeon to Bellevue Hospital; Consulting Surgeon to Bureau of Surgical and Medical Relief for the Out-Door Poor, at Bellevue Hospital; to the Central Dispensary; and to the Hospital for the Ruptured and Crippled; Fellow of the New York Academy of Medicine, etc.

THE PRINCIPLES AND PRACTICE OF SURGERY. Illustrated with four hundred and sixty-seven engravings on wood. Royal 8vo, 954 pages. In muslin, \$5.50; or in leather, \$6.

**Helferich, H., M.D.,**

Professor at the University of Greifswald.

**ATLAS OF TRAUMATIC FRACTURES AND LUXATIONS.**  
With a Brief Treatise. One 12mo volume, containing 64 full-page plates, embracing 106 figures, finely executed by chromo-lithography, with complete descriptive text. Muslin, \$3.00 *net*. (Volume III. of Wood's Medical Hand Atlases.)

**Herman, G. Ernest, M.B. Lond., F.R.C.P.,**

Senior Obstetric Physician to the London Hospital; Physician to the General Lying-in Hospital; President of the Obstetrical Society of London; Examiner in Midwifery to the Royal College of Surgeons; Late Physician to the Royal Maternity Charity.

**DIFFICULT LABOR: A Guide to its Management for Students and Practitioners.** 460 pages, demi-octavo, including complete index, muslin, \$2.25.

**Holden, Luther, M.D.,**

Ex-President and Member of the Court of Examiners of the Royal College of Surgeons of England; Consulting Surgeon to Saint Bartholomew's and the Foundling Hospitals; assisted by

**Shuter, James, F.R.C.S., M.A., M.B. Cantab.,**

Assistant Surgeon to the Royal Free Hospital; late Demonstrator of Physiology, and Assistant Demonstrator of Anatomy, at Saint Bartholomew's Hospital.

**HUMAN OSTEOLOGY.** Comprising a Description of the Bones, with Delineations of the Attachments of the Muscles, the General and Microscopic Structure of Bone and its Development. Sixth edition. With sixty-six full-page lithographic plates, and eighty-nine wood-engravings. 8vo, 285 pages, muslin, \$1.00.

**Household Practice.**

See WOOD'S HOUSEHOLD PRACTICE.

**Hudson, E. D., Jr., A.M., M.D.,**

Professor of General Medicine and Diseases of the Chest in the New York Polyclinic; Physician to Bellevue Hospital, etc.

**A MANUAL OF THE PHYSICAL DIAGNOSIS OF THORACIC DISEASES.** 8vo, 162 pages, profusely illustrated. Muslin, \$1.50.

**Hutchinson, Jonathan, F.R.S.**

**THE PEDIGREE OF DISEASE.** Being Six Lectures on Temperament, Idiosyncrasy, and Diathesis. Muslin, \$1.25.

## **Jakob, Dr. Christfried,**

Practising Physician in Bamberg; formerly First Assistant in the Medical Clinic at Erlangen.  
With an Introduction by

## **Prof. Dr. Ad. v. Strümpell,**

Translated and Edited (authorized) by

## **Joseph Collins, M.D.,**

Instructor of Nervous and Mental Diseases, New York Post-Graduate Medical School, etc.

**AN ATLAS OF THE NERVOUS SYSTEMS, NORMAL AND PATHOLOGICAL**, together with a Sketch of the Anatomy, Pathology, and Therapy of the same. One 12mo volume, containing 78 plates, embracing 118 figures, superbly executed by chromo-lithography, with complete descriptive text and an introductory chapter. The plates are executed most exquisitely, and form a collection most unique in neurological literature. Muslin, \$3.00 *net*. (Volume II. of Wood's Medical Hand Atlases.)

## **Ingals, E. Fletcher, A.M., M.D.,**

Professor of Laryngology and Practice of Medicine, Rush Medical College; Professor of Diseases of Throat and Chest, Northwestern University Women's Medical School; Professor of Laryngology and Rhinology, Chicago Polyclinic, etc.

**LECTURES ON THE DIAGNOSIS AND TREATMENT OF DISEASES OF THE CHEST, THROAT, AND NASAL CAVITIES.** Including Physical Diagnosis and Diseases of the Lungs, Heart, and Aorta; Laryngology and Diseases of the Pharynx, Larynx, Nose, Thyroid Gland, and Esophagus. Third edition, revised and enlarged. 8vo, 708 pages, 240 illustrations, including colored plate of stained tubercle bacilli. Muslin, \$5.00.

## **Kaposi, Dr. Moriz,**

Professor of Dermatology and Syphilis, and Chief of the Clinic and Division for Skin Diseases in the Vienna University.

**PATHOLOGY AND TREATMENT OF DISEASES OF THE SKIN.** For Practitioners and Students. Translation of the latest German edition. 8vo, 684 pages, 84 illustrations, and a colored plate, muslin, \$5.50; leather, \$6.50.

## **Kellogg, Theodore H., A.M., M.D.,**

New York, Late Superintendent, Willard State Hospital; former Physician-in-Chief of the New York City Asylum for the Insane, etc., etc.

**A TEXT-BOOK ON MENTAL DISEASES**, for the Use of Students and Practitioners of Medicine. One large octavo volume, of 792 pages, illustrated by engravings and charts. Muslin, \$6.00.



**Kirchhoff, Dr. Theodore,**

Physician to the Schleswig Insane Asylum and Privat-Docent at the University of Kiel.

**HANDBOOK OF INSANITY FOR PRACTITIONERS AND STUDENTS.** Illustrated with eleven plates. 8vo, 362 pages. Muslin, \$2.75 ; flexible leather, gilt top, \$3.50.

**Kirkes' Handbook of Physiology.**

**HANDBOOK OF PHYSIOLOGY.** By W. MORRANT BAKER, F.R.C.S., and VINCENT DORMER HARRIS, M.D. Lond., F.R.C.P. American Revision of Chapters on the Nervous System by CHARLES L. DANA, A.M., M.D., Professor of Nervous and Mental Diseases in the New York Post-Graduate Medical School, and in Dartmouth Medical College ; Visiting Physician to Bellevue Hospital ; Neurologist to the Montefiore Home ; ex-President of the American Neurological Association, etc.

One volume of 860 pages, 8vo, illustrated with a colored plate and five hundred and sixteen illustrations, muslin, \$4.00 ; leather, \$5.00

**Knies, Max, M.D.,**

Professor Extraordinary at the University of Freiburg.

**THE EYE AND ITS DISEASES, IN RELATION TO THE DISEASES OF OTHER ORGANS.** Translated and edited by H. D. NOYES, M.D. 8vo, 470 pages, illustrated, muslin, \$4.25.

**Landau, Prof. Dr. Leopold, and  
Landau, Dr. Theodor,**

Berlin.

**THE HISTORY AND TECHNIQUE OF THE VAGINAL RADICAL OPERATION.** Translated by B. L. EASTMAN, M.D., Berlin, and ARTHUR E. GILES, M.D., London. One volume, octavo, with numerous original illustrations. Muslin, \$2.00 *net*.

**Lehman, Prof. K. B., and  
Neumann, Rudolf,**

Würzburg.

**ATLAS AND ESSENTIALS OF BACTERIOLOGY.** Sixty-three superbly executed chromo-lithographic plates, with description facing each, and 150 pages of text, comprising an introduction to the Morphology of Bacteria, a chapter on technique, index, etc. Muslin, \$3.00 *net*. (Volume VI. of Wood's Medical Hand Atlases.)

**Ling, P. He.**

**SYSTEM OF MANUAL TREATMENT AS APPLICABLE TO SURGERY AND MEDICINE.** By ARVID KELLGREN, M.D. Edin 8vo, 151 pages, with 79 illustrations, muslin, \$1.25.



**Liveing, Robert, A.M. and M.D. Cantab., F.R.C.P. Lond.**

Lecturer on Dermatology to the Middlesex Hospital Medical School; lately Physician to the Middlesex Hospital; Author of "Notes on the Treatment of Skin Diseases," "Elephantiasis Græcorum," etc.

A HANDBOOK ON THE DIAGNOSIS OF SKIN DISEASES.  
One volume, 16mo, 266 pages, muslin, \$1.50.

NOTES ON THE TREATMENT OF SKIN DISEASES. One  
volume, 16mo, 127 pages, muslin, \$1.00.

**Loomis, Alfred L., M.D., LL.D.,**

Professor of Pathology and Practical Medicine, in the Medical Department of the University of the City of New York; Visiting Physician to Bellevue Hospital, etc.

A TEXT-BOOK OF PRACTICAL MEDICINE. One handsome 8vo  
volume of 1,147 pages, illustrated by two hundred and eleven engravings.  
Eleventh edition. Muslin, \$6.00; leather, \$7.00.

LESSONS IN PHYSICAL DIAGNOSIS. Tenth revised edition. One  
volume, 8vo, 290 pages, illustrated, muslin, \$3.00.

**Macfarlane, A. W., M.D.,**

Fellow of the Royal College of Physicians, Edinburgh; Fellow of the Royal Medical and Chirurgical Society of London; Examiner in Medical Jurisprudence in the University of Glasgow, etc.

INSOMNIA AND ITS THERAPEUTICS. 8vo, 302 pages, muslin,  
\$1.75.

**Macnaughton-Jones, H., M.D., M.Ch.**

PRACTICAL MANUAL OF DISEASES OF WOMEN AND UTER-  
INE THERAPEUTICS, for Students and Practitioners. Seventh  
revised and enlarged edition. One volume of 933 pages, small 8vo,  
illustrated by 565 wood-engravings. Muslin, \$5.00.

**Matthieu, A.,**

Physician to the Paris Hospitals.

TREATMENT OF THE DISEASES OF THE STOMACH AND  
INTESTINES. 8vo, 285 pages, muslin, \$2.50; flexible leather, gilt  
top, round corners, \$3.25.

**May, Charles H., M.D.,**

Instructor in Ophthalmology, New York Polyclinic; and

**Mason, Charles F., M.D.,**

Late Assistant Surgeon, U.S.A.

AN INDEX OF MATERIA MEDICA. With Prescription Writing,  
including Practical Exercises. 32mo, muslin, \$1.00. (Wood's Pocket  
Manuals.)

**Mauthner, Ludwig,**

Royal Professor of the University of Vienna.

THE SYMPATHETIC DISEASES OF THE EYE. Translated  
from the German by WARREN WEBSTER, M.D., JAMES A. SPAULDING,  
M.D. 12mo, 220 pages, muslin, \$2.00.

**McGillicuddy, T. J., A.M., M.D.**

FUNCTIONAL DISORDERS OF THE NERVOUS SYSTEM IN WOMEN. One volume, 8vo, uniform with the Medical Practitioners' Library, 373 pages, illustrated by forty-five wood-engravings and two chromo-lithographic plates. Extra muslin, \$3.00; flexible leather, \$3.50.

**Medical Record.**

A WEEKLY JOURNAL OF MEDICINE AND SURGERY. Subscription price, \$5.00 per year.

**Medical Record Visiting List.**

See VISITING LIST.

**Millard, H. B., M.D.**

A TREATISE ON BRIGHT'S DISEASE OF THE KIDNEYS; ITS PATHOLOGY, DIAGNOSIS, AND TREATMENT. Third edition, revised and enlarged. 8vo, 322 pages, numerous original illustrations, muslin, \$3.00.

**Miller, Maurice N., M.D.**

PRACTICAL MICROSCOPY. A Course of Normal Histology for Students and Practitioners of Medicine. 8vo, 217 pages, muslin, \$2.00.

**Moore, John William.**

A TEXT-BOOK OF THE ERUPTIVE AND CONTINUED FEVERS. 8vo, 535 pages, illustrated with lithographic plates and temperature charts, muslin, \$4.00.

**Morris, Henry, M.A., M.B. Lond., F.R.C.S.,**

Surgeon to and Lecturer on Surgery at the Middlesex Hospital; Member of the Council and of the Court of Examiners of the Royal College of Surgeons, England; Examiner in Surgery in the University of London.

INJURIES AND DISEASES OF THE GENITAL AND URINARY ORGANS. One volume of 494 pages, 8vo, illustrated by 96 wood engravings, muslin, \$4.00.

**Morrow, P. A., A.M., M.D.,**

Clinical Professor of Venereal Diseases; Consulting Surgeon to the Bellevue Out-Door Department, etc.

VENEREAL MEMORANDA. A Manual for the Student and Practitioner. Second edition. 32mo, muslin, \$1.00. (Wood's Pocket Manuals.)

ATLAS OF SKIN AND VENEREAL DISEASES. One volume, half morocco. \$25.00. (Subscription.)

DRUG ERUPTIONS. A Clinical Study of the Irritant Effect of Drugs upon the Skin. 8vo, 206 pages, one lithographed plate, muslin, \$1.75.

**Moullin, C. W. Mansell.**

SPRAINS, THEIR CONSEQUENCES AND TREATMENT. 8vo  
221 pages, muslin, \$1.50.

**Murrell, William, M.D., F.R.C.P. Lond.**

A MANUAL OF MATERIA MEDICA AND THERAPEUTICS.  
By Special Arrangement with the Author; Revised to Conform with  
American Practice by FREDERICK A. CASTLE, M.D., New York. One  
volume of 522 pages, 8vo, with complete index. Muslin, \$4.00.

CLINICAL LECTURES ON THE PREVENTION OF CONSUMPTION. 12mo, muslin, \$1.00.

**Noman, Dr. D. Van Haren.**

Professeur e. o. de clinique dermatologique et syphiligraphique à la Faculté de Médecine d'Amsterdam.

CASUISTIQUE ET DIAGNOSTIC PHOTOGRAPHIQUE DES  
MALADIES DE LA PEAU. Consisting of photographic plates,  
with descriptive text. This work is to be in ten parts, unbound.  
Price complete, \$20.00. (Six parts are published so far.)

**Noyes, Henry D., M.D.,**

Professor of Ophthalmology and Otology in Bellevue Hospital Medical College; Executive Surgeon to the New York Eye and Ear Infirmary; recently President of the American Ophthalmological Society, etc.

A TEXT-BOOK ON DISEASES OF THE EYE. Royal 8vo, 832  
pages, richly illustrated with chromo-lithographic plates and 269  
engravings. Second Edition. Muslin, \$6.00; sheep, \$7.00.

**Obstetrics and Gynecology.**

CYCLOPÆDIA OF OBSTETRICS AND GYNECOLOGY. Twelve  
volumes, 8vo. Profusely illustrated by colored plates. Lithographs  
in tint and nearly two thousand wood-engravings. The set, \$12.00.

**Oldberg, Oscar, Ph.D.,**

Member of the Committee of Revision of the Pharmacopœia of the United States; Author of the "Unofficial Pharmacopœia," "The Metric System in Medicine," etc.; formerly Medical Purveyor of the United States Marine Hospital Service; and Professor of Materia Medica in the National College of Pharmacy, Washington, D. C., etc.; and

**Wall, Otto A., M.D., Ph.G.,**

Professor of Materia Medica, Therapeutics, and Pharmacy in the Missouri Medical College, and of Materia Medica and Botany in the St. Louis College of Pharmacy; Member of the Committee of Revision of the Pharmacopœia of the United States, etc.

A COMPANION TO THE UNITED STATES PHARMACOPŒIA.  
Second edition. Being a Commentary on the Pharmacopœia, and containing the Descriptions, Properties, Uses, and Doses of all Official and numerous Unofficial Drugs and Preparations in current use in the United States, together with Practical Hints, Working Formulas, etc., designed as a ready reference book for Pharmacists, Physicians, and Students, with over 300 original illustrations. Muslin, \$5.00; leather, \$6.00.

**Parkes, E., M.D.**

A MANUAL OF PRACTICAL HYGIENE. Edited by F. S. B. FRANÇOIS DE CHAUMONT, M.D. Sixth edition. With an Appendix, giving the American practice in matters relating to hygiene. Prepared by and under the supervision of FREDERICK N. OWEN, Civil and Sanitary Engineer. Two volumes in one, 8vo, 946 pages. Illustrated with nine full-page plates and fine wood-engravings, muslin, \$5.00.

**Partridge, Edward L., M.D.,**

New York City.

THE OBSTETRICAL REMEMBRANCER. Profusely illustrated with miniature wood-engravings. (Wood's Pocket Manuals.) 32mo, muslin, \$1.00.

**Paschkis, Heinrich.**

COSMETICS. A Treatise for Physicians. A complete translation from the German edition. 204 pages, paper, 50c.

**Peabody, George L., M.D.,**

New York.

SUPPLEMENT TO ZIEMSEN'S CYCLOPÆDIA OF THE PRACTICE OF MEDICINE. By Various Authors. In one royal 8vo volume, 844 pages, bound to correspond. Muslin, \$6.00; leather, \$7.00; half morocco, \$8.00; also extra muslin (not corresponding), \$6.00; and red leather, \$7.00.

**Pictures for Physicians' Offices and Libraries.**

Edward Jenner, the first Inoculation of Vaccine, May 14, 1796.  
Andrew Vesalius, the Anatomist.  
Spoonful Every Hour.  
The Sick Wife.  
Ambrose Paré Demonstrating the Use of Ligatures.  
The Young Mother.  
The Village Doctor.

Prof. Charcot's Clinic at the "Salpêtrière" Hospital, Before the Operation.  
The Rebellious Patient.  
Study in Anatomy.  
William Harvey Demonstrating the Circulation of the Blood.  
The Anatomical Lecture.  
The Accident.  
The Doctor.  
Anæsthesia.

Size of each, 19x24 inches. Price, each \$1.00. Illustrated catalogue sent upon application.

Prof. Billroth's Clinic, Vienna, size 24x32, \$2.00.

**Piffard, Henry G., A.M., M.D.,**

Professor of Dermatology, University of the City of New York; Surgeon to the Charity Hospital, etc.

A GUIDE TO URINARY ANALYSIS FOR THE USE OF PHYSICIANS AND STUDENTS. 8vo, 88 pages, illustrated, \$1.25.



**Porter, William Henry, M.D.,**

Late Professor of Clinical Medicine and Pathology in the New York Post-Graduate Medical School and Hospital; Curator to the Presbyterian Hospital.

A PRACTICAL TREATISE ON RENAL DISEASES AND URINARY ANALYSIS. 360 pages, one hundred illustrations, muslin, \$3.00.

**Pozzi, S., M.D.**

Professeur Agrégé à la Faculté de Médecine, Chirurgien de l'Hôpital Lourcine-Pascal, Paris.

TREATISE ON MEDICAL AND SURGICAL GYNÆCOLOGY.

Translated from the third French edition, under the supervision of BROOKS H. WELLS, M.D., Lecturer on Gynæcology at the New York Polyclinic; Fellow of the New York Obstetrical Society and the New York Academy of Medicine. One royal 8vo volume of about 936 pages, illustrated by 600 fine wood-engravings. Muslin, \$5.50 *net*; leather, \$6.50 *net*.

**Rabagliati, A., M.A., F.R.C.S. Ed.,**

Honorary Gynecologist, Late Senior Honorary Surgeon, Bradford Infirmary.

SYMPTOMS WHICH SIMULATE DISEASE OF THE PELVIC ORGANS IN WOMEN, and their treatment by Allo-Piesto-Myo-Kinetics (Massage), and by Auto-Piesto-Myo-Kinetics (Self-movements of muscles under pressure). 8vo, illustrated by six full-page heliogravure plates, muslin, \$1.50.

**Reference Handbook of the Medical Sciences.**

By various authors. Edited by ALBERT H. BUCK, M.D., Clinical Professor of the Diseases of the Ear, in the College of Physicians and Surgeons, New York; Consulting Aural Surgeon, New York Eye and Ear Infirmary. Nine volumes, imperial 8vo, muslin, \$6.00 per volume; leather, \$7.00 per volume; half-morocco, \$8.00 per volume. (Subscription.) Circular on application.

**Reynolds, Edward,**

Fellow of the American Gynecological Society; of the Obstetric Society of Boston, etc.; Assistant in Obstetrics in Harvard University; Physician to Out-Patients of the Boston Lying-In Hospital, etc.

PRACTICAL MIDWIFERY. A HANDBOOK OF TREATMENT. Third Revised Edition. 8vo, 427 pages, small octavo, 121 illustrations. Muslin, \$2.50.

**Ringer, Sidney, M.D.,**

Professor of the Principles and Practice of Medicine in University College; Physician to University College Hospital.

A HANDBOOK OF THERAPEUTICS, WITH DIETARY AND INDEX OF DISEASES. Twelfth edition. 8vo, 535 pages, muslin \$5.00; leather, \$6.00.



**Robson, A. W. Mayo, F.R.C.S.,**

Leeds, Eng.

**DISEASES OF THE GALL-BLADDER AND BILE DUCTS.**One volume, octavo. Muslin, \$2.00 *net*.**Rockwell, A. D., A.M., M.D.****THE MEDICAL AND SURGICAL USES OF ELECTRICITY.**

Entirely rewritten from the former book by Beard and Rockwell. One large 8vo volume of 628 pages, profusely illustrated. Muslin, \$4.50; sheep, \$5.50.

**Roosa, D. B. St. John, M.D.,**

Professor of Diseases of the Eye and Ear in the University of the City of New York; Surgeon to the Manhattan Eye and Ear Hospital; Consulting Surgeon to the Brooklyn Eye and Ear Hospital; formerly President of the Medical Society of the State of New York; Corresponding Member of the Medico-Chirurgical Society of Edinburgh; Member of the Medical Society of the County of New York, etc.

**TEXT-BOOK ON DISEASES OF THE EYE.** Including a sketch of its anatomy. Illustrated by 178 engravings and 2 chromo-lithographic plates. Muslin, \$5.50; leather, \$6.50.**A PRACTICAL TREATISE ON THE DISEASES OF THE EAR, INCLUDING THE ANATOMY OF THE ORGAN.** Seventh edition. One volume, 8vo, 763 pages. Illustrated by 140 wood-engravings and chromo-lithographs, muslin, \$5.50; leather, \$6.50.**A VEST-POCKET MEDICAL LEXICON.** Being a Dictionary of the Words, Terms, and Symbols of Medical Science. Collated from the best authorities, with the additions of words not before introduced into a Lexicon. With an Appendix. Third revised and enlarged edition. One volume, 64mo, roan, 75c., or tucks, \$1.00.**THE OLD HOSPITAL, AND OTHER PAPERS.** Being the second revised and enlarged edition of "A Doctor's Suggestions." 8vo, 320 pages, gilt top, uncut, dark olive cloth, \$3.00.**Roosa, D. B. St. John, M.D., and Ely, Edward T., M.D.****OPHTHALMIC AND OTIC MEMORANDA.** (Wood's Pocket Manuals.) Fourth edition. One volume, 32mo, 298 pages, muslin, \$1.00.**Roth, Otto.****THE MATERIA MEDICA OF MODERN MEDICINE.** Second edition. Translated from the revised German edition and adapted to the U. S. Pharmacopœia. 8vo, 467 pages, muslin, \$2.00.

**Sachs, B., M.D.,**

Professor of Mental and Nervous Diseases in the New York Polyclinic; Consulting Neurologist to the Mt. Sinai Hospital; Neurologist to the Montefiore Home for Chronic Invalids; Ex-President of the American Neurological Association.

A TREATISE ON THE NERVOUS DISEASES OF CHILDREN.  
For Physicians and Students. 8vo, 688 pages, profusely illustrated with colored plate, muslin, \$5.00.

**Salomonsen, C. J., and Trelease, William.**

BACTERIOLOGICAL TECHNOLOGY FOR PHYSICIANS. Authorized translation from the Second Revised Danish edition. 8vo, 163 pages, 72 illustrations, muslin, \$1.25.

**Schaeffer, Oscar, M.D. Heidelberg.**

ANATOMICAL ATLAS OF OBSTETRIC DIAGNOSIS AND TREATMENT. Sixty-four beautifully executed full-page chromolithographic plates, containing 142 figures. Together with 250 pages of descriptive text and treatise. (Volume IV. of Wood's Medical Hand Atlases.) Muslin, \$3.00 *net*.

ATLAS AND ELEMENTS OF GYNÆCOLOGY. Sixty-four superbly executed colored plates, comprising 159 figures, with description and a short treatise, illustrated by 54 wood engravings. (Volume V. of Wood's Medical Hand Atlases.) Muslin, \$3.00 *net*.

**Schmidt-Rimpler, Dr. Herman,**

Professor of Ophthalmology and Diseases of the Ophthalmoscopic Clinic at Marburg, Germany.

OPHTHALMOLOGY AND OPHTHALMOSCOPY. A Complete Treatise upon Diseases and Injuries to the Eye, for Students and Practitioners of Medicine. Revised and edited by D. B. ST. JOHN ROOSA, M.D., Professor of Diseases of the Eye and Ear in the New York Post-Graduate Medical School; Surgeon to the Manhattan Eye and Ear Hospital, etc. Royal 8vo, 571 pages, illustrated by 183 wood-engravings and by three colored plates, muslin, \$6.00.

**Schreiber, August.**

GENERAL ORTHOPEDICS, INCLUDING ORTHOPEDIC SURGERY. Complete translation from the original German edition. 8vo, 357 pages, 388 illustrations, muslin, \$1.75.

**Schroeder, Aimée Raymond, M.D.**

HEALTH NOTES FOR YOUNG WIVES. 12mo, 218 pages, fancy half cloth, \$1.00.

**Seguin, E., M.D.**

MEDICAL THERMOMETRY AND HUMAN TEMPERATURE.  
8vo, 445 pages, illustrated with diagrams, muslin, \$3.50.

**Semeleder, Dr. Friedrich,**

Physician in Ordinary to his Majesty, the Emperor of Mexico; Member of the Royal Medical Society of Vienna and of the Medical Society of Pantheon in Paris; Formerly Member of the Medical Faculty of the University of Vienna, and Surgeon to the Branch Hospital at Gumpendorf.

**RHINOSCOPY AND LARYNGOSCOPY: THEIR VALUE IN PRACTICAL MEDICINE.** Translated from the German by EDWARD T. CASWELL, M.D. With woodcuts and two chromo-lithographic plates. 8vo, 191 pages, muslin, \$3.25.

**Sexton, Samuel, M.D.,**

Aural Surgeon to the New York Eye and Ear Infirmary; Fellow of the American Otological Society; Fellow of the New York Academy of Medicine; Member of the Medical Society of the County of New York, and of the Practitioners' Society of New York.

**THE EAR AND ITS DISEASES, BEING PRACTICAL CONTRIBUTIONS TO THE STUDY OF OTOTOLOGY.** Edited by CHRISTOPHER J. COLLES, M.D. 8vo, 473 pages, illustrated, muslin, \$4.00.

**Skene, Alexander J. C., M.D.,**

Professor of the Diseases of Women, in the Long Island College Hospital; Fellow of the American Gynecological Society; Corresponding Member of the Gynecological Society of the County of Kings, and of the Obstetrical Society of New York.

**DISEASES OF THE BLADDER AND URETHRA IN WOMEN.** Second revised edition. 8vo, 374 pages, illustrated, muslin, \$3.00.

**Smart, Chas., M.D., Major U.S.A.**

**A HANDBOOK FOR THE HOSPITAL CORPS OF THE UNITED STATES ARMY AND STATE MILITARY FORCES.** Nearly 600 pages, measuring about 4 x 5 inches, green leather, with metal clasp, \$2.50.

**Starr, M. Allen, M.D., Ph.D.,**

Professor of Diseases of the Mind and Nervous System; College of Physicians and Surgeons, New York.

**FAMILIAR FORMS OF NERVOUS DISEASE.** With illustrations, diagrams, and charts. 8vo, 350 pages, muslin, \$3.00.

**BRAIN SURGERY.** 8vo, 306 pages, illustrated, muslin, \$3.00.

**Steel, J. H., M.D.**

**OUTLINE OF EQUINE ANATOMY.** A Manual for the use of Veterinary Students in the Dissecting Room. 12mo, 312 pages, muslin, \$3.00.

**Stewart, R. W., M.D., M.R.C.S.,**

Pittsburg.

**THE DISEASES OF THE MALE URETHRA.** One volume of 229 pages, post-octavo, illustrated by numerous wood-engravings. Muslin, \$2.50; flexible morocco, \$3.25. (*Medical Practitioners' Library.*)

## **Sternberg, George M., M.D., F.R.M.S.,**

Surgeon-General U. S. Army ; Director of the Hoagland Laboratory, Brooklyn, N. Y. ; Honorary Member of the Epidemiological Society of London, of the Royal Academy of Medicine of Rome, of the Academy of Medicine of Rio de Janeiro, of the American Academy of Medicine, etc.

A TEXT-BOOK OF BACTERIOLOGY. One volume, large 8vo, 693 pages, illustrated by heliotype and chromo-lithographic plates and two hundred engravings in black and colors. Extra muslin, \$5.50 ; brown sheep, \$6.50.

A MANUAL OF BACTERIOLOGY. Illustrated by heliotype and chromo-lithographic plates and 321 engravings in black and colors. 8vo, 850 pages, muslin, \$8.00 *net* ; leather, \$9.00 *net*.

IMMUNITY: PROTECTIVE INOCULATIONS IN INFECTIOUS DISEASES AND SERUM-THERAPY. Post 8vo, 332 pages, muslin, \$2.50 ; flexible morocco, \$3.25.

## **Stewart, T. Grainger, M.D.,**

Fellow of the Royal College of Physicians ; Physician to the Royal Infirmary ; Lecturer on Clinical Medicine ; formerly Pathologist to the Royal Infirmary ; Lecturer in General Pathology at Surgeons' Hall, and Physician to the Royal Hospital for Sick Children ; Extraordinary Member and formerly President of the Royal Medical Society of Edinburgh.

CLINICAL LECTURES ON ALBUMINURIA. 8vo, 261 pages, muslin, \$2.25.

A PRACTICAL TREATISE ON BRIGHT'S DISEASE OF THE KIDNEYS. 8vo, 334 pages, illustrated with seven lithographic plates, muslin, \$4.50.

## **Supplement to the International Encyclopedia of Surgery.**

One imp. 8vo volume, of 1,136 pages, illustrated. Muslin, \$7.50 *net* ; leather, \$8.50 *net* ; half morocco, \$9.50 *net*. (To subscribers to the set : muslin, \$6.00 ; leather, \$7.00 ; half morocco, \$8.00.) Circular on application.

## **Surgery, International Encyclopedia of.**

By Various Authors. Edited by Dr. JOHN ASHHURST, Jr. Six volumes, imp. 8vo, of about 950 pages each, muslin, \$6.00 ; leather, \$7.00 ; half morocco, \$8.00. Send for circulars. (Subscription.)

## **Thomas, John J.**

Revised and Enlarged by WILLIAM H. S. WOOD, Esq.

THE AMERICAN FRUIT CULTURIST. Twentieth edition. Post 8vo, 784 pages, illustrated by 800 wood-engravings. Muslin, \$2.50.



## Twentieth Century Practice.

AN INTERNATIONAL ENCYCLOPEDIA OF MODERN MEDICAL SCIENCE. By Leading Authorities of Europe and America. Edited by THOMAS L. STEDMAN, M.D., New York City. To be completed in 20 volumes, royal 8vo, published one every three months. Muslin, \$5.00, leather, \$6.00; half morocco, \$7.50. Twelve volumes now ready. (Subscription.) Circulars on application.

## Visiting List (Medical Record), or Physician's Diary.

Containing, besides the diary, much useful information on many subjects of daily interest to the physician. Prices: For thirty patients a week; handsome red or black leather binding, with or without dates, \$1.25, for sixty patients a week, same style, with or without dates, \$1.50. Removable, fitting into black sealskin and calf wallets, from \$2.50 to \$4.00. Send for a circular.

## Walsham, W. J., M.B., C.M. Aberd., F.R.C.S. Eng.,

Senior Assistant-Surgeon, Surgeon-in-Charge of the Orthopedic Department and Lecturer in Anatomy, St. Bartholomew's Hospital, London.

## Hughes, Wm. Kent, M.B. Lond., M.B. Melb., M.R.C.S. Eng., L.R.C.P. Lond.,

Orthopedic Surgeon, St. Vincent's Hospital; Assistant Surgeon Children's Hospital, Melbourne.

THE DEFORMITIES OF THE HUMAN FOOT, WITH THEIR TREATMENT. 558 pages, post 8vo, profusely illustrated by 296 engravings, muslin, \$4.50.

## Warren, J. Collins, M.D.,

Assistant Professor of Surgery, Harvard University; Surgeon to the Massachusetts General Hospital; Member American Surgical Association; Honorary Fellow Philadelphia Academy of Surgery.

THE HEALING OF ARTERIES AFTER LIGATURE IN MAN AND ANIMALS. 8vo, 184 pages. Superbly illustrated with twelve full-page plates in black and colors. Muslin, \$3.25.

## Whittaker, J. T., M.D., LL.D.,

Professor of the Theory and Practice of Medicine, Medical College of Ohio, etc., etc.

A PRACTICE OF MEDICINE, PREPARED FOR STUDENTS AND PRACTITIONERS. 8vo, 700 pages, illustrated, muslin, \$5.75; leather, \$6.50.

## Wendt, Edmund C., M.D.,

Curator of St. Francis Hospital; Pathologist and Curator of the New York Infant Hospital, etc.

A TREATISE ON CHOLERA. Edited and prepared in Association with JOHN C. PETERS, M.D., New York; JOHN B. HAMILTON, M.D., Surgeon-General U. S. Marine Hospital Service, and ELY McCLELLAN, M.D., Surgeon U. S. Army. 8vo, 503 pages, illustrated with maps and engravings, muslin, \$3.00.



**Williams, J. W. Hume,**

Of the Middle Temple, Barrister-at-Law, London.

UN SOUNDNESS OF MIND IN ITS LEGAL AND MEDICAL CONSIDERATIONS. A complete reprint of this important work. 8vo, 166 pages, muslin, \$1.50.

**Williams, W. Roger, F.R.C.S. Lond.**

A MONOGRAPH ON DISEASES OF THE BREAST; THEIR PATHOLOGY AND TREATMENT, WITH SPECIAL REFERENCE TO CANCER. One volume of 580 pages, 8vo, illustrated by 76 wood-engravings. Muslin, \$4.50.

**Witthaus, R. A., A.M., M.D.,**

Professor of Medical Chemistry and Toxicology in the University of Vermont; Member of the Chemical Societies of Paris and Berlin, etc.

THE MEDICAL STUDENT'S MANUAL OF CHEMISTRY. (American Series of Medical Text-Books.) Fourth revised edition. 556 pages and 62 woodcuts, muslin, \$3.75.

ESSENTIALS OF CHEMISTRY AND TOXICOLOGY. For the Use of Students in Medicine. Twelfth edition. (Wood's Pocket Manuals.) 32mo, 319 pages, muslin, \$1.00.

GUIDE TO URINALYSIS AND TOXICOLOGY. For Students and Practitioners. Third revised edition. Oblong 12mo, 153 pages, interleaved, muslin, \$1.00.

**Witthaus, R. A., M.D., and Becker, T. C., Esq.**

With a staff of Collaborators.

MEDICAL JURISPRUDENCE AND TOXICOLOGY. Four volumes, 8vo, bound in muslin and leather, at \$5.00 and \$6.00 respectively. (Subscription.) Circulars on application.

**Wood's Household Practice of Medicine, Hygiene, and Surgery.**

A Practical Treatise for the Use of Families, Travellers, Seamen, Miners, and others. By Various Authors. 8vo, 765 pages, illustrated by colored lithographic plates and five hundred fine wood-engravings. Muslin, \$5.00.

**Wood's Index Rerum.**

The finest arrangement yet devised for all ready record and reference purposes. For professional use in recording your cases, or in grouping your cases from your case books. Patent index. Vowel arrangement. Bound for permanency in ledger binding, \$5.00.

**Wood's Pocket Lexicon.**

See ROOSA.

**Woodburn, W. D., L.D.S.,**

Glasgow, Scotland.

ON EXTRACTION, WITH NOTES ON THE ANATOMY AND PHYSIOLOGY OF THE TEETH. One volume, 12mo, 104 pages, profusely illustrated. Extra muslin, \$1.50.

**Wyeth, John A., M.D.,**

Member of the New York County Medical Society, the New York Pathological Society; Honorary Member of the College of Physicians and Surgeons of Little Rock, Arkansas.

A HANDBOOK OF MEDICAL AND SURGICAL REFERENCE. 12mo, 279 pages, tucks, \$1.50.

ESSAYS IN SURGICAL ANATOMY AND SURGERY. 8vo, 262 pages, illustrated, muslin, \$2.00.

**Yonge, Eugene S., M.B.**

Master in Surgery, University of Edinburg, etc.

AIDS TO SURGICAL ANATOMY. 16 mo, 158 pages, \$1.00.

**Ziegler, Ernst,**


Professor of Pathological Anatomy and of General Pathology in the University of Freiburg.

TEXT-BOOK OF GENERAL PATHOLOGY. Royal 8vo, 610 pages, with 456 illustrations in black and numerous tints, and a chromo-lithographic plate. Eighth revised and enlarged edition. Muslin, \$5.50; leather, \$6.50.

**Ziemssen, H. Von, M.D. Munich.**

CYCLOPÆDIA OF THE PRACTICE OF MEDICINE. By Various Authors. Complete in twenty volumes, royal 8vo. Price per volume, in muslin, \$5.00; in leather, \$6.00; in half morocco, \$7.50 (Subscription.) Circulars on application.


## A SELECTION OF



PICTURES


.. FOR .. PHYSICIANS' OFFICES

Send for complete catalogue of same. A heliotype sheet of the series, in miniature, will be sent upon receipt of four cents in stamps













LIBRARY OF CONGRESS



0 022 194 775 7